

EHRHART POLYNOMIAL FOR LATTICE SQUARES, CUBES AND HYPERCUBES

EUGEN J. IONASCU

Communicated by Alexandru Zaharescu

In this paper, we are constructing integer lattice squares, cubes or hypercubes in \mathbb{R}^n with $n \in \{2, 3, 4\}$. We find a complete description of their Ehrhart polynomial. We characterize all the integer squares in \mathbb{R}^4 , in terms of two Pythagorean quadruple representations of the form $a^2 + b^2 + c^2 = d^2$, and then prove a parametrization in terms of two quaternions of all such squares. We introduce the sequence of *almost perfect squares in dimension n* . In dimension two, this is very close to the sequence A194154 (in OEIS).

AMS 2010 Subject Classification: 52C07, 05A15, 68R05.

Key words: Ehrhart polynomial, linear Diophantine equations, lattice square, lattice cube, quaternions, icube, Pythagorean quadruple, twin vectors.

1. INTRODUCTION

Eugène Ehrhart [9, 10] proved that given a d -dimensional compact simplicial complex in \mathbb{R}^n ($1 \leq d \leq n$), denoted here generically by \mathcal{P} , whose vertices are in the lattice \mathbb{Z}^n , there exists a polynomial $L(\mathcal{P}, t) \in \mathbb{Q}[t]$ of degree d , associated with \mathcal{P} , satisfying

$$L(\mathcal{P}, t) = \text{the cardinality of } \{t\mathcal{P}\} \cap \mathbb{Z}^n, \quad t \in \mathbb{N}.$$

It is very interesting that one can say more about three of the coefficients of $L(\mathcal{P}, t)$:

$$(1) \quad L(\mathcal{P}, t) = \text{Vol}(\mathcal{P})t^d + \frac{1}{2}\text{Vol}(\partial\mathcal{P})t^{d-1} + \dots + \chi(\mathcal{P}),$$

where $\text{Vol}(\mathcal{P})$ is the usual volume of \mathcal{P} normalized with respect to the sublattice containing \mathcal{P} , $\text{Vol}(\partial\mathcal{P})$ is the surface area of \mathcal{P} normalized with respect to the sublattice on each face of \mathcal{P} and $\chi(\mathcal{P})$ is the Euler characteristic of \mathcal{P} (in the sense of polytopal complexes, so for a convex polytope it is equal to one). In general, the other coefficients in (1) may have complicated expressions in terms of the vertices of \mathcal{P} .

It is also known that the number of points in the interior of $t\mathcal{P}$ is given by $(-1)^d L(\mathcal{P}, -t)$. For a polytope that is a cross product of two polytopes, the Ehrhart polynomial is the product of the corresponding smaller degree Ehrhart polynomials. In this article, we are studying this polynomial for lattice squares, cubes, and hypercubes in \mathbb{R}^n . Such objects have been constructed and counted in several works (see [13, 15, 20, 25]).

2. SQUARES

2.1. Squares in two dimensions

How general can a lattice square in two dimensions look like? We may assume that one of the vertices is at the origin. Then the other vertices, in counterclockwise order, are given by $A(a, b)$, $B(a-b, a+b)$ and $C(-b, a)$ where a and b are integers, not both zero, and such that $\gcd(a, b) = 1$. Using the facts mentioned in the Introduction, the polynomial in (1) is simply $E_{\square}(t) = (a^2 + b^2)t^2 + 2t + 1$ and of course, $E_{\square}(t) = (a^2 + b^2)t^2 - 2t + 1$, $t \in \mathbb{N}$.




Fig. 1 – Square with $a = 5$ and $b = 2$.

In Figure 1, we see the 28 points inside of the square $OABC$, with $A(5, 2)$, $B(3, 8)$ and $C(-2, 5)$. One interesting problem here is to determine the sequence

of possible lattice points in the interior of such a square. The first hundred terms of this sequence, listed in increasing order, are included in the table below:

0	1	4	5	9	12	13	16	17	24	25	28	33	36	37	40	41	49	52	57
60	61	64	65	72	73	81	84	85	88	96	97	100	101	105	108	112	113	116	121
124	129	133	136	144	145	148	153	156	161	168	169	172	177	180	181	184	192	193	196
197	201	204	209	217	220	221	225	228	229	232	240	241	249	256	257	264	265	268	273
276	280	288	289	292	293	297	301	304	305	312	313	316	324	325	328	336	337	345	348

Let us call this sequence the *almost perfect squares* sequence. In Figure 2, we see some of the squares that show that the above numbers are indeed in the sequence. As a curiosity, 2015 is not in this sequence but 2016 is. This sequence is very close to A194154 in the OEIS (On-Line Encyclopedia of Integer Sequences), but 20 is the first term that is not in our sequence.




Fig. 2 – Various squares.

Let us give a reformulation of the exact form of the Ehrhart polynomial above in conjunction with the reciprocity property (about interior points), which can be shown independently using Pick's Theorem.

PROPOSITION 2.1. *Given a, b with $\gcd(a, b) = 1$ and $t \in \mathbb{N}$, then*

$$\#\{(x, y) \in \mathbb{Z}^2 : ax + by, ay - bx \in [1, t(a^2 + b^2) - 1]\} = (a^2 + b^2)t^2 - 2t + 1.$$

Proof. Counting the points inside the square $OABC$, with $O(0, 0)$, $A(a, b)$, $B(a - b, a + b)$ and $C(-b, a)$, is equivalent to count the solutions (x, y) of the system $(x, y) = \alpha(ta, tb) + \beta(-tb, ta)$ with $\alpha, \beta \in (0, 1)$. Since $\alpha = (x, y) \cdot (a, b)/t(a^2 + b^2) = \frac{ax + by}{t(a^2 + b^2)}$ and $\beta = (x, y) \cdot (-b, a)/t(a^2 + b^2) = \frac{ay - bx}{t(a^2 + b^2)}$ we see that the constraints on α and β is equivalent to $ax + by, ay - bx \in [1, t(a^2 + b^2) - 1]$. Hence, the result follows from the Ehrhart polynomial expression. \square

So, every term of the sequence of almost perfect squares is the answer to a counting as in Proposition 2.1, which in particular gives the justification of one of our earlier claims:

$$\sharp\{(x, y) \in \mathbb{Z}^2 : 44x + 9y, 44y - 9x \in [1, 2016]\} = 2016.$$

2.2. Squares in three dimensions

In \mathbb{R}^3 , we can obtain a lattice square by taking two orthogonal vectors with integer coordinates of the same length: $u = (a, b, c)$, $v = (a', b', c')$ such that $a^2 + b^2 + c^2 = a'^2 + b'^2 + c'^2 = \ell$ and $aa' + bb' + cc' = 0$. In [13], two such vectors are called *twin* vectors. The number of such twin vectors having a given length is calculated in [13]. In [20], this concept is generalized to m dimensions and called m -icube (a set of m vectors in \mathbb{Z}^n , mutually orthogonal and of the same length). Since we are interested in generalizing these ideas to any dimension, we are going to use this terminology also.

Although it does not make much of a difference it is natural to assume that the square is *irreducible*, i.e., $\gcd(a, b, c, a', b', c') = 1$. If we set $d = \gcd(a, b, c)$, $d' = \gcd(a', b', c')$ and $D = \gcd(bc' - b'c, ac' - a'c, ab' - b'a)$, we have the following more general formula, than in two dimensions, for the $E_{\square}(t)$.

THEOREM 2.2. *The Ehrhart polynomial of a lattice square embedded into \mathbb{R}^3 , described above and with the notation introduced is given by*

$$(2) \quad E_{\square}(t) = Dt^2 + (d + d')t + 1.$$

Proof. The fundamental domain of the lattice containing the triangle has a volume equal to $\sqrt{n_1^2 + n_2^2 + n_3^2}$ where $\vec{n} = [n_1, n_2, n_3]$ is a vector normal to the plane containing the triangle and such that $\gcd(n_1, n_2, n_3) = 1$ (see [1] and [19]). Such a vector is given clearly by the cross-product of u and v : $n = \frac{1}{D}u \times v = \frac{1}{D}(bc' - b'c, -(ac' - a'c), ab' - b'a)$. Let us observe that because of Lagrange's Identity $\ell^2 = (a^2 + b^2 + c^2)(a'^2 + b'^2 + c'^2) = (aa' + bb' + cc')^2 + (bc' - b'c)^2 + (ac' - a'c)^2 + (ab' - b'a)^2$, we conclude that $|n| = \frac{\ell}{D}$. Hence, we see that the first coefficient of the Ehrhart polynomial is $\sqrt{\ell^2}/(\frac{\ell}{D}) = D$. Similar argument goes for the second coefficient. \square

Standard Examples. Let us analyze some examples which are not xy , xz or yz -plane examples. First, let us take $u = (3, -3, 0)$ and $v = (1, 1, 4)$; observe that $|u|^2 = |v|^2 = 18$ and $u \cdot v = 0$. In this case $n = 6(-2, -2, 1)$, and the equation of the plane is $2x + 2y - z = 0$ and $E_{\square}(t) = 6t^2 + 4t + 1$. We know that $E_{\square}(-1) = 3$ represents the number of lattice points in the interior of the square. We notice that 3 (and $E_{\square}(-2) = 17$) was not an almost perfect

number. Hence we may want to generalize that sequence to *almost perfect squares in dimension n* .

What is interesting is that in the plane $2x + 2y - z = 0$ we have another square: $\bar{u} = (2, -1, 2)$ and $\bar{v} = (-1, 2, 2)$. Also, the square above can be written in terms of this one basically like in two dimensions: $u = \bar{u} - \bar{v}$ and $v = \bar{u} + \bar{v}$. For this square, $n = 3(-2, -2, 1)$ and so the Ehrhart polynomial is $E_{\{\bar{u}, \bar{v}\}}(t) = 3t^2 + 2t + 1$. We see that $E_{\{\bar{u}, \bar{v}\}}(-1) = 2$ which is yet new for the almost perfect square sequence compared with dimension two.

If we take the twin vectors $u = (6, -2, 3)$ and $v = (-2, 3, 6)$ we get $n = 7(-3, -6, 2)$ and so the Ehrhart polynomial is $E(t) = 7t^2 + 2t + 1$. There seem to be very few numbers that are not perfect squares in 3D: 7, 14, 23,...

For an example of an irreducible square, for which d and d' in Theorem 2.2 are both greater than one, we refer to $u = 5(8, 12, 9)$ and $v = 17(0, -3, 4)$. Its polynomial is $E(t) = (5t + 1)(17t + 1)$.

How does one construct such squares in \mathbb{R}^3 ? We observed in the proof of Theorem 2.2 that every such square is contained in a plane whose normal $n = (n_1, n_2, n_3)$ satisfies

$$(3) \quad \ell^2 = n_1^2 + n_2^2 + n_3^2.$$

A solution of this type of Diophantine equation is usually referred in the literature as a *Pythagorean quadruple*. For the number of primitive solutions of (3) in terms of ℓ , we refer the reader to a recent paper of Werner Hürlimann [15] but also [4]. There are at least two other terms used for these quadruples of integers: cuboids [15] and *Lorenz quadruples* (see [14]). We include in the next tables all primitive solutions for odd values $\ell \in \{1, 3, ..19\}$:

ℓ	[a,b,c], gcd(a,b,c) = 1, c even	ℓ	[a,b,c], gcd(a,b,c) = 1, c even
1	[1,0,0]	11	[9, 2, 6], [7, 6, 6]
3	[1,2,2]	13	[5, 12, 0], [3, 4, 12]
5	[3,4,0]	15	[5, 14, 2], [11, 2, 10]
7	[3,6,2]	17	[15, 0, 8], [1, 12, 12], [9, 12, 8]
9	[1, 4, 8], [7, 4, 4]	19	[1, 6, 18], [17, 6, 6], [15, 6, 10]

So, having a Pythagorean quadruple, how can we construct a related twin set of vectors? We include a partial answer to this question at this point, but we will include the complete solution later in this paper.

THEOREM 2.3. *Suppose that $u' = (a, b, c)$ satisfies $n_1a + n_2b + n_3c = 0$, where $n_1^2 + n_2^2 + n_3^2 = \ell^2$ with all variables involved being integers. Then there exist $v = (a', b', c')$ such that $\ell u'$ and v define a lattice square in the plane of normal $n = (n_1, n_2, n_3)$.*

Proof. We define v to be the cross-product of u' and n . Clearly v is perpendicular to n and so it is in the right plane. Since u' and n are perpendicular, $|v| = |u'| |n| = \ell |u'|$. \square

It is natural to look for the “smallest” square in the plane $n_1a + n_2b + n_3c = 0$. This leads us to the shortest vector problem (SVP), *i.e.*, finding a non-zero vector in a lattice of minimum norm. An interesting problem, at this point, is to characterize all the values ℓ so that $\sqrt{\ell}$ appears as side-lengths for an lattice embedded square in \mathbb{R}^m , $m \geq 2$. If we denote this set by \mathcal{L} , from the two dimensions construction, we see that \mathcal{L} is invariant under multiplication with numbers which are sums of two squares. For $m = 3$, it is also clear all the numbers of the form $4^k(8s - 1)$ are not in \mathcal{L} , since these are not representable as sums of three squares (Legendre’s three-square theorem). Also, 3, 11 or 19 (see the sequence A223732 in OEIS) are not in \mathcal{L} , since they have only one representation as sum of three squares and these representations contain only odd numbers. Hence the sum $aa' + bb' + cc'$ is also odd and so, it cannot be zero. We will describe the set \mathcal{L} at the end of the subsection. We will see that in \mathbb{R}^4 and beyond, $\mathcal{L} = \mathbb{N}$.

We have the following parametrization for those squares whose side-lengths are natural numbers of the form $x^2 + y^2 + z^2 + t^2$ (see [18] and [25]):

$$(4) \quad u := (2ty + 2zx, 2tz - 2yx, t^2 - z^2 - y^2 + x^2), \quad v := (2zy - 2tx, \\ z^2 - t^2 + x^2 - y^2, 2tz + 2yx),$$

having normal vector $n = (-x^2 + t^2 - y^2 + z^2, -2(tx + zy), 2(ty - zx))$, $|n| = x^2 + y^2 + z^2 + t^2$. Next, we show that, as in [18] and [25], the following similar parametrization for the solutions of (3) takes place. The proof is essentially the same as in [25] but we include it for the convenience of the reader.

THEOREM 2.4 ([25]). *Every primitive solution of (3), after a permutation of variables and change of signs, is given by $n_1 = 2(zy - tx)$, $n_2 = 2(tz + yx)$ and $n_3 = z^2 - t^2 + x^2 - y^2$, and $\ell = x^2 + y^2 + z^2 + t^2$ for some integers x, y, z and t .*

Proof. As usual, a primitive solution of (3) is one for which $\gcd(n_1, n_2, n_3) = 1$. We must have ℓ odd since otherwise $\gcd(n_1, n_2, n_3) \geq 2$. In this case, one of the n_i must be odd and the other two even. Without loss of generality, let us assume that n_1 and n_2 are even. Then, we have

$$(\ell - n_3)(\ell + n_3) = n_1^2 + n_2^2.$$

We know that $\ell + n_3 = 2\alpha$ and $\ell - n_3 = 2\beta$ are both even, and then the above equality becomes

$$\alpha\beta = (n_1/2)^2 + (n_2/2)^2 = A^2 + B^2 = (A + Bi)(A - Bi),$$

where $A = n_1/2$ and $B = n_2/2$. A Gaussian prime of the form $p = 4k + 3$ which divides $\alpha\beta$, must divide both A and B . So, it cannot divide both α and β because it then divides $\ell = (\alpha + \beta)/2$ and $n_3 = (\alpha - \beta)/2$, which implies that $\gcd(n_1, n_2, n_3) \geq p$. Hence, α and β are both sums of two squares. If we let $\gcd(A + iB, \alpha) = t + iy$ and $\gcd(A + iB, \beta) = x + iz$. Let us make the observation that these equalities are defined up to a unit, *i.e.*, ± 1 or $\pm i$.

Since α is real, $t - iy$ divides α and so $\alpha = (t^2 + y^2)\alpha'$. But every prime factor of α , appears as a factor of either $A + Bi$ or $A - Bi$. Taking into account the multiplicities we see that α divides $t^2 + y^2$ and so $\alpha' = 1$. Similarly, we have $\beta = x^2 + z^2$. Also, $t + iy$ divides $A + iB$ and also $x + iz$ divides $A + iB$. Hence $A + iB = (t + iy)(x + iz)k$ and from here $A^2 + B^2 = (t^2 + y^2)(x^2 + z^2)|k|^2$ which together with what we have shown earlier forces $|k| = 1$. So, by changing x and y we can assume that $k = 1$. Hence we have $A = tx - zy$ and $B = xy + tz$. Then $\ell = \alpha + \beta = x^2 + y^2 + z^2 + t^2$ and $n_3 = \alpha - \beta = t^2 + y^2 - x^2 - z^2$. \square

Theorem 2.4 allows us to characterize \mathcal{L} . The result is not new, as we found recently, it is contained in [13]. We include a proof of it, based on our development of the ideas.

THEOREM 2.5. *The set of all ℓ so that $\sqrt{\ell}$ is the side-lengths for an embedded square in \mathbb{Z}^3 is the set of positive integers which are sums of two squares.*

Proof. In one direction, *i.e.*, \mathcal{L} contains the set of all positive integers which are sums of two squares, we can observe that the lengths for squares in two dimensions are of the form $\sqrt{a^2 + b^2}$. For the other direction, if we start with an arbitrary square, \mathcal{S} , in \mathbb{Z}^3 , let us suppose the side is $\sqrt{\ell}$. As in the proof of Theorem 2.3, this square is contained in a plane of normal $n = (n_1, n_2, n_3)$, with $n_1^2 + n_2^2 + n_3^2 = \ell^2$. By Theorem 2.4, we can find x, y, z and t , such that $n_1 = 2zy - 2tx$, $n_2 = 2tz + 2yx$ and $n_3 = z^2 - t^2 + x^2 - y^2$. Then the parametrization (4) gives a square in the same plane as \mathcal{S} , whose sides are ℓ . We can then write the vector of this square in the basis given by two of the vectors in \mathcal{S} : $w = \alpha u + \beta v$ for some α and $\beta \in \mathbb{Q}$. Then, taking norms we get $\ell^2 = \alpha^2 \ell + \beta^2 \ell$ so $\ell = \alpha^2 + \beta^2$. This shows that $\ell = A^2 + B^2$ for some $A, B \in \mathbb{Z}$. \square

Theorem 2.3 and Theorem 2.5 imply the next consequence which one can also prove elementary.

COROLLARY 2.6. *Given a point $P = (x, y, z) \in \mathbb{Z}^3$ in the plane of equation $n_1x + n_2y + n_3z = 0$ with $n_1^2 + n_2^2 + n_3^2 = \ell^2$ for some $\ell \in \mathbb{N}$, then the number $x^2 + y^2 + z^2$ is actually a sum of two squares.*

2.3. Squares in \mathbb{R}^4

Perhaps, one of the simplest ways to construct squares in \mathbb{Z}^4 is to take two squares in two dimensions and “add” them together, each on its own dimensions. In other words, the two vectors u and v that define the square, as we have seen before, are of the form $u = (a, b, c, d)$ and $v = (-b, a, -d, c)$. We have clearly $u \cdot v = 0$ and $\|u\|^2 = \|v\|^2 = a^2 + b^2 + c^2 + d^2$. This simple example allows us to answer the question about the possible side-lengths of such squares. By Lagrange’s Four Square Theorem, we see that every natural number is a possible side-length of a square. Are these squares the most general situation that one can expect? We observe that u and v are essentially the first two rows of the following *pseudo-orthogonal matrix* (every two rows are mutually orthogonal and have the same norm):

$$O_{a,b,c,d} = \begin{bmatrix} a & b & c & d \\ -b & a & d & -c \\ -c & -d & a & b \\ -d & c & -b & a \end{bmatrix}.$$

It is not difficult to see that the product of matrices like these ($O^T O = tI$) is of the same form. This allows one to define a certain multiplication on vectors in \mathbb{R}^4 which is exactly the quaternion multiplication that we will be using latter.

Example 1. Let us compute the Ehrhart polynomial for a particular case like this which gives a new value for its sides. We have $u = (2, 1, 1, 1)$ and $v = (-1, 2, -1, 1)$. Using the same idea as in the three dimensions we are computing two normal vectors that define the orthogonal space of the plane generated by u and v . For a generic point $P = (x, y, z, t) \in \mathbb{R}^4$, the two equations, computed with the help of the cross-product in the three dimensions, are

$$3x - y - 5z = 0 \text{ and } 2y + z - 3t = 0.$$

We observe that both u and v satisfy these equations. Since these two equations can be solved for y and t , we can find two integer vectors which generate the lattice \mathbb{Z}^4 intersected with the plane defined by u and v : $y = 3x - 5z$ and $t = 2x - 3z$ which gives the two vectors $\alpha = (1, 3, 0, 2)$ and $\beta = (0, -5, 1, -3)$. We know then that these two vectors form a fundamental domain, so we need to compute the area of the parallelogram generated by these two vectors: $A = |\alpha||\beta|\sin(\gamma)$ where $\cos \gamma = \alpha \cdot \beta / |\alpha||\beta| = -3/\sqrt{10}$. This gives $A = 7$ and so, the Ehrhart polynomial is

$$E_{\square}(t) = t^2 + 2t + 1 = (t + 1)^2.$$

For a set of vectors \mathcal{S} in \mathbb{R}^4 we denote as usual by \mathcal{S}^\perp the set of all vectors $x \in \mathbb{R}^4$ perpendicular to every vector v in \mathcal{S} , i.e.,

$$\mathcal{S}^\perp = \{(x_1, x_2, x_3, x_4) \in \mathbb{R}^4 \mid x_1v_1 + x_2v_2 + x_3v_3 + x_4v_4 = 0 \text{ for all } v = (v_1, v_2, v_3, v_4), v \in \mathcal{S}\}.$$

THEOREM 2.7. (i) *Given integer vectors $u = (u_1, u_2, u_3, u_4)$ and $v = (v_1, v_2, v_3, v_4)$ such that*

$$0 < \ell = u_1^2 + u_2^2 + u_3^2 + u_4^2 = v_1^2 + v_2^2 + v_3^2 + v_4^2, \quad u_1v_1 + u_2v_2 + u_3v_3 + u_4v_4 = 0,$$

then there exist an odd $k \in \mathbb{N}$ dividing ℓ and two vectors w_1 and w_2 with integer coordinates such that $w_1 = (0, \alpha_1 - \beta_1, \alpha_2 - \beta_2, \alpha_3 - \beta_3)$ and $w_2 = (\alpha_3 - \beta_3, -\alpha_2 - \beta_2, \alpha_1 + \beta_1, 0)$ with

$$(5) \quad k^2 = \alpha_1^2 + \alpha_2^2 + \alpha_3^2 = \beta_1^2 + \beta_2^2 + \beta_3^2,$$

α_i and β_i of the same parity and $u, v \in \{w_1, w_2\}^\perp$. One can permute the coordinates of u and v and/or change their signs in order to have w_1 and w_2 linearly independent.

(ii) *If $\gcd(\alpha_1, \beta_1, \alpha_2, \beta_2, \alpha_3, \beta_3) = 1$ then the volume of the fundamental domain of the minimal lattice containing u and v is equal to k .*

Proof. Using Lagrange's Identity, we get

$$\ell^2 = \left(\sum_{i=1}^4 u_i^2\right)\left(\sum_{i=1}^4 v_i^2\right) = \left(\sum_{i=1}^4 u_i v_i\right)^2 + \sum_{1 \leq i < j \leq 4} (u_i v_j - u_j v_i)^2.$$

Since $u_1v_1 + u_2v_2 + u_3v_3 + u_4v_4 = 0$, this implies that

$$(6) \quad \ell^2 = \sum_{1 \leq i < j \leq 4} (u_i v_j - u_j v_i)^2.$$

If we denote by $\Delta_{ij} = (-1)^{i-j}(u_i v_j - u_j v_i)$, it is not difficult to check that

$$(7) \quad \Delta_{12}\Delta_{34} - \Delta_{13}\Delta_{24} + \Delta_{14}\Delta_{23} = 0.$$

This helps us change (6) into

$$(8) \quad \ell^2 = (\Delta_{12} \pm \Delta_{34})^2 + (\Delta_{13} \mp \Delta_{24})^2 + (\Delta_{14} \pm \Delta_{23})^2.$$

One can check that

$$(9) \quad \begin{cases} u_1(0) + u_2\Delta_{34} + u_3\Delta_{24} + u_4\Delta_{23} = 0 \\ v_1(0) + v_2\Delta_{34} + v_3\Delta_{24} + v_4\Delta_{23} = 0 \\ u_1\Delta_{23} + u_2\Delta_{13} + u_3\Delta_{12} + u_4(0) = 0 \\ v_1\Delta_{23} + v_2\Delta_{13} + v_3\Delta_{12} + v_4(0) = 0. \end{cases}$$

Let us observe that if 2 divides ℓ , then all Δ_{ij} are divisible by 2, because all u_i (v_i) are all even or all odd. In order to prove the claim, we simplify (8) by the greatest power of 2 possible and take $\alpha_1 = (\Delta_{12} + \Delta_{34})/\ell'$, $\alpha_2 = (-\Delta_{13} + \Delta_{24})/\ell'$, $\alpha_3 = (\Delta_{14} + \Delta_{23})/\ell'$, $\beta_1 = (\Delta_{12} - \Delta_{34})/\ell'$, $\beta_2 = (-\Delta_{13} - \Delta_{24})/\ell'$, and $\beta_3 = (\Delta_{14} - \Delta_{23})/\ell'$, with ℓ' so that $\gcd(\alpha_1, \beta_1, \alpha_2, \beta_2, \alpha_3, \beta_3) = 1$.

It is easy to check that if $\alpha_3 \neq \beta_3$ then the two vectors w_1 and w_2 are linearly independent and so the orthogonal space $\{w_1, w_2\}^\perp$ is two-dimensional. By permuting coordinates and/or changing their signs, we can insure that $\alpha_3 - \beta_3 = 2\Delta_{23}/\ell' \neq 0$, since by (6) and the assumption that $\ell > 0$, we see that not all of the Δ_{ij} are equal to zero.

(ii) Let us denote by \mathcal{L} the minimal lattice which contains the given square $\{u, v\}$ and by V its volume. As we observed in part (i) this lattice is the same as $\mathbb{Z}^4 \cap \{w_1, w_2\}^\perp$ if $\alpha_3 \neq \beta_3$. Let us observe that the two vectors $U := (0, \Delta_{12}, -\Delta_{13}, \Delta_{14})$ and $V := (\Delta_{14}, -\Delta_{24}, \Delta_{34}, 0)$ are in \mathcal{L} . This shows that $2U/\ell' = (0, \alpha_1 + \beta_1, \alpha_2 + \beta_2, \alpha_3 + \beta_3)$ and $2V/\ell' = (\alpha_3 + \beta_3, -\alpha_2 + \beta_2, \alpha_1 - \beta_1, 0)$ are in \mathcal{L} also. The area of the parallelogram determined by these two vectors is given by the square root of the Gramian determinant

$$\begin{aligned} & \begin{vmatrix} (\alpha_1 + \beta_1)^2 + (\alpha_2 + \beta_2)^2 + (\alpha_3 + \beta_3)^2 & (\alpha_1 + \beta_1)(\beta_2 - \alpha_2) + (\alpha_2 + \beta_2)(\alpha_1 - \beta_1) \\ (\alpha_1 + \beta_1)(\beta_2 - \alpha_2) + (\alpha_2 + \beta_2)(\alpha_1 - \beta_1) & (\alpha_3 + \beta_3)^2 + (\alpha_2 - \beta_2)^2 + (\alpha_1 - \beta_1)^2 \end{vmatrix} = \\ & = \begin{vmatrix} 2(k^2 + \alpha_1\beta_1 + \alpha_2\beta_2 + \alpha_3\beta_3) & 2(\alpha_1\beta_2 - \beta_1\alpha_2) \\ 2(\alpha_1\beta_2 - \beta_1\alpha_2) & 2(k^2 + \alpha_3\beta_3 - \alpha_2\beta_2 - \alpha_1\beta_1) \end{vmatrix}. \end{aligned}$$

This means that the vectors U/ℓ' and V/ℓ' , still in \mathcal{L} , form a parallelogram of an area, which is the square root of

$$\begin{aligned} & (1/4)[(k^2 + \alpha_3\beta_3)^2 - (\alpha_2\beta_2 + \alpha_1\beta_1)^2 - (\alpha_1\beta_2 - \beta_1\alpha_2)^2] = \\ & (1/4)[k^4 + 2k^2\alpha_3\beta_3 + \alpha_3^2\beta_3^2 - (k^2 - \alpha_3^2)(k^2 - \beta_3^2)] = k^2((\alpha_3 + \beta_3)/2)^2. \end{aligned}$$

It follows that V divides $k|\alpha_3 + \beta_3|/2$. In a similar way, we can show that V divides $k|\alpha_i \pm \beta_i|/2$ for $i = 1, 2, 3$. Given the assumption that $\gcd(\alpha_1, \beta_1, \alpha_2, \beta_2, \alpha_3, \beta_3) = 1$, we conclude that V divides k .

In order to conclude that $V = k$, let us look at the Gram determinant of the vectors $w_1/(\alpha_3 - \beta_3)$ and $w_2/(\alpha_3 - \beta_3)$, which if added to \mathcal{L} extends the lattice and gives a basis for it. Since its volume can be determined by the same method as above, a similar calculation gives $V' = k/|\alpha_3 - \beta_3|$. This implies that V/V' is an integer, or k divides $V|\alpha_3 - \beta_3|$. Since we can obtain similarly that k divides $V|\alpha_i \pm \beta_i|$, we conclude that $V = k$. \square

Example 2. As in the case of equilateral triangles (see [17]), there is a converse of Theorem 2.7. Let us take the first odd ℓ for which we get two essentially different representations as in (5):

$$11^2 = 9^2 + 6^2 + 2^2 = 7^2 + 6^2 + 6^2.$$

We can then choose $\alpha_1 = 9$, $\beta_1 = 7$, $\alpha_2 = 6$, $\beta_2 = 6$, $\alpha_3 = 2$, and $\beta_3 = 6$. Then the two vectors, defined in Theorem 2.7, are $w_1 = (0, 2, 0, -4)$ and $w_2 = (-4, -12, 16, 0)$. Then the space $\{w_1, w_2\}^\perp$ is defined by the equations $y - 2t = 0$ and $x + 3y - 4z = 0$. These can be simplified to $y = 2t$ and $x = 4z - 6t$. Because the generic vector in $\{w_1, w_2\}^\perp \cap \mathbb{Z}^4$ is $u = (4z - 6t, 2t, z, t) = z(4, 0, 1, 0) + t(-6, 2, 0, 1)$ we can calculate, as before the “volume” of the fundamental domain and obtain indeed 11. Then we get a quadratic form $QF(z, t) = (4z - 6t)^2 + (2t)^2 + z^2 + t^2 = 17z^2 - 48zt + 41t^2$ that should lead to the solutions we need solving the Diophantine equation $QF(z, t) = 11k$. It turns out that the smallest multiple for which we have solutions is $k = 11$ and then two of the vectors which define the square are $u = (-4, 8, 5, 4)$ and $v = (10, 2, 4, 1)$. Its Ehrhart polynomial is then $E_\square(t) = 11t^2 + 2t + 1$. We observe that we get new terms, such as 10, 94, and 266 for instance, for the sequence of almost perfect squares in dimension 4, compared to dimension 2. This is an example when all the Δ_{ij} are divisible by 11.

As we have seen in [17], the following result gives a certain converse to Theorem 2.7.

THEOREM 2.8. *Given k odd, and two different representations*

$$k^2 = a^2 + b^2 + c^2 = a'^2 + b'^2 + c'^2, \text{ with } \gcd(a, b, c, a', b', c') = 1, \quad c' > c,$$

and a, a' both odd. Then if we set $\Delta_{12} = \frac{a'-a}{2}$, $\Delta_{34} = \frac{a+a'}{2}$, $\Delta_{13} = -\frac{b'-b}{2}$, $\Delta_{24} = \frac{b+b'}{2}$, $\Delta_{14} = \frac{c+c'}{2}$, and $\Delta_{23} = \frac{c'-c}{2}$ the equations (7) and (8) are satisfied. Moreover, the two dimensional space \mathcal{S} of all vectors $[u, v, w, t] \in \mathbb{Z}^4$, such that

$$(10) \quad \begin{cases} (0)u + \Delta_{34}v + \Delta_{24}w + \Delta_{23}t = 0 \\ \Delta_{23}u + \Delta_{13}v + \Delta_{12}w + (0)t = 0 \end{cases}$$

contains a family of squares.

Proof. By construction we see that (8) is true. Since a, a' are odd, the others must be even and so, all numbers defined above are integers. We have then indeed

$$\Delta_{12}\Delta_{34} - \Delta_{13}\Delta_{24} + \Delta_{14}\Delta_{23} = (1/4)[a'^2 - a^2 - (b^2 - b'^2) + (c'^2 - c^2)] = 0.$$

One can also check that $k^2 = \sum_{i < j} \Delta_{ij}^2$. We see that the assumption $\Delta_{23} > 0$ insures that the equations (10) define a two dimensional space in \mathbb{R}^4 . We can solve the equations (10) for t and u :

$$t = -\frac{\Delta_{34}v + \Delta_{24}w}{\Delta_{23}}, \quad u = -\frac{\Delta_{13}v + \Delta_{12}w}{\Delta_{23}}, \quad \text{with } v, w \in \mathbb{Z}.$$

If we denote a generic point $P \in \mathbb{R}^4$ in the plane (10), in terms of v and w , *i.e.*,

$$P(v, w) = \left[-\frac{\Delta_{13}v + \Delta_{12}w}{\Delta_{23}}, v, w, -\frac{\Delta_{34}v + \Delta_{24}w}{\Delta_{23}} \right],$$

we observe that for two pairs of the parameters, (v, w) and (v', w') , we obtain $\Delta'_{23} = wv' - vw'$,

$$\Delta'_{12} = v \left(-\frac{\Delta_{13}v' + \Delta_{12}w'}{\Delta_{23}} \right) + \frac{\Delta_{13}v + \Delta_{12}w}{\Delta_{23}} v' = \Delta_{12}\Delta'_{23}/\Delta_{23},$$

and similarly $\Delta'_{13} = \Delta_{13}\Delta'_{23}/\Delta_{23}$, etc.

We want to show that a non-zero square of the form $OP(v, w)P(v', w')Q$ ($OQ = OP(v, w) + OP(v', w')$) exists for some integer values of v, w, v' and w' . We need to have $OP(v, w)^2 = OP(v', w')^2 = \ell$ and $P(v, w) \cdot P(v', w') = 0$.

If such a square exists, by Theorem 2.7, we may want ℓ of the form $\ell = k\ell'$ for some $\ell' \in \mathbb{N}$.

By Lagrange's identity, calculations similar to those in the derivation of (6), show that if $|P(v, w)|^2 = |P(v', w')|^2 = k\ell'$ then

$$(P(v, w) \cdot P(v', w'))^2 = k^2\ell'^2 - \left(\sum_{i,j} \Delta_{ij}^2 \right) \left(\frac{\Delta'_{23}}{\Delta_{23}} \right)^2 = k^2\ell'^2 - k^2 \left(\frac{\Delta'_{23}}{\Delta_{23}} \right)^2.$$

Hence, to get $P(v, w) \cdot P(v', w') = 0$, it is enough to have $\Delta'_{23} = \ell'\Delta_{23}$ (provided that we have already shown that $|P(v, w)|^2 = |P(v', w')|^2 = k\ell'$). In other words, a square exists if there exist integer pairs (v, w) and (v', w') such that

$$(11) \quad |OP(v, w)|^2 = |OP(v', w')|^2 = k \frac{\Delta'_{23}}{\Delta_{23}}.$$

The essential object in this analysis is the quadratic form

$$QF(v, w) := |OP(u, v)|^2 = \left(\frac{\Delta_{34}v + \Delta_{24}w}{\Delta_{23}} \right)^2 + v^2 + w^2 + \left(\frac{\Delta_{13}v + \Delta_{12}w}{\Delta_{23}} \right)^2,$$

or

$$QF(v, w) = \frac{(\Delta_{34}^2 + \Delta_{13}^2 + \Delta_{23}^2)v^2 + 2(\Delta_{34}\Delta_{24} + \Delta_{13}\Delta_{12})vw + (\Delta_{24}^2 + \Delta_{12}^2 + \Delta_{23}^2)w^2}{\Delta_{23}^2}.$$

We use Lagrange's identity again, which we will write in the form

$$(\alpha^2 + \beta^2 + \gamma^2)(\zeta^2 + \eta^2 + \theta^2) = (\alpha\zeta - \beta\eta + \gamma\theta)^2 + (\alpha\eta + \beta\zeta)^2 + (\alpha\theta - \gamma\zeta)^2 + (\beta\theta + \gamma\eta)^2.$$

Then the determinant of the form $QF(v, w)$ (excluding its denominator) is equal to

$$\begin{aligned} -\Delta/4 &= (\Delta_{34}^2 + \Delta_{13}^2 + \Delta_{23}^2)(\Delta_{12}^2 + \Delta_{24}^2 + \Delta_{23}^2) - (\Delta_{34}\Delta_{24} + \Delta_{13}\Delta_{12})^2 = \\ &= (\Delta_{12}\Delta_{34} - \Delta_{13}\Delta_{24} + \Delta_{23}^2)^2 + (\Delta_{12}\Delta_{23} - \Delta_{34}\Delta_{23})^2 + (\Delta_{13}\Delta_{23} + \Delta_{24}\Delta_{23})^2 \Rightarrow \\ -\Delta/4 &= \Delta_{23}^2 [(\Delta_{23} - \Delta_{14})^2 + (\Delta_{12} - \Delta_{34})^2 + (\Delta_{13} + \Delta_{24})^2] = \Delta_{23}^2(k^2) \\ &\Rightarrow \Delta = -(2k\Delta_{23})^2. \end{aligned}$$

This implies that for $v_0 = -(\Delta_{34}\Delta_{24} + \Delta_{13}\Delta_{12})$ and $w_0 = \Delta_{13}^2 + \Delta_{23}^2 + \Delta_{34}^2$ we get

$$QF(v_0, w_0) = k^2(\Delta_{34}^2 + \Delta_{13}^2 + \Delta_{23}^2),$$

and

$$(12) \quad QF(v, w) = \frac{(w_0v - v_0w)^2 + k^2w^2\Delta_{23}^2}{\Delta_{23}^2w_0}.$$

So, by (11) and (12) we need to have solutions (v, w) and (v', w') of

$$(13) \quad (w_0v - v_0w)^2 + k^2w^2\Delta_{23}^2 = (w_0v' - v_0w')^2 + k^2w'^2\Delta_{23}^2 = k\Delta_{23}w_0(wv' - vw').$$

In general, if the Diophantine quadratic equation $x^2 + y^2 = n$ has a solution (x, y) , then it has other solutions such as $(y, -x)$.

Given (v, w) , let us assume that (v', w') gives exactly this other solution $(x = w_0v - v_0w, y = kw\Delta_{23})$ when substituted in (13):

$$\begin{cases} w_0v' - v_0w' = kw\Delta_{23} \\ k\Delta_{23}w' = -(w_0v - v_0w). \end{cases}$$

This allows us to solve for v' and w' in order to calculate Δ'_{23} :

$$v' = \frac{v_0^2w - v_0w_0v + k^2\Delta_{23}^2w}{2k\Delta_{23}w_0}, \quad w' = \frac{v_0w - w_0v}{k\Delta_{23}},$$

$$\Delta'_{23} = wv' - vw' = \frac{(w_0v - v_0w)^2 + k^2w^2\Delta_{23}^2}{k\Delta_{23}w_0}.$$

This means that (13) is automatically satisfied with this choice of (v', w') . In fact, we have shown that, for every integer values of (v, w) , such that

$$(14) \quad (w_0v - v_0w)^2 + k^2w^2\Delta_{23}^2 = (kw_0\ell')\Delta_{23}^2$$

taking (v', w') as above, we automatically get a square determined by vectors $OP(v, w), P(v', w') \in \mathbb{Q}^4$, which may not have integer coordinates. In order for (14) to have rational solutions, in v and w , we need to have $kw_0\ell'$ a positive integer which is a product of primes of the form $4s+1$, and all the other primes, in its prime factorization, have even exponents. Clearly, there exists a smallest positive integer ℓ' with this property. \square

As we have seen in Proposition 2.2 [17], we can similarly show the existence in every plane of equations (10), of a square that is *minimal*, in the sense that the side-lengths are the smallest possible. Then, a similar result to Proposition 2.2 [17] takes place: *every square in the same plane can be written in terms of a square that is minimal (same formulae as in 2 dimensions)*. For a proof, we just invite the reader to study Figure 3 (a proof “without words” exercise).




Fig. 3 –Minimal squares.

As in dimension 3, we also have parametric formulae for squares, which are more general than the ones at the beginning of the subsection. The two vectors are

(15)

$$u = \pm(-ta-zb+yc-xd, -tb+za-yd-xc, -tc-zd-ya+xb, -td+zc+yb+xa)$$

$$v = \pm(ax-by-cz-dt, ay+bx+ct-dz, az-bt+cx+dy, at+bz-cy+dx)$$

and one can check that $u \cdot v = 0$ and $|u|^2 = |v|^2 = (x^2 + y^2 + z^2 + t^2)(a^2 + b^2 + c^2 + d^2)$. The two relations as in (5) can be simplified by $(x^2 + y^2 + z^2 + t^2)^2$ and $(a^2 + b^2 + c^2 + d^2)^2$ respectively and reduced to

(16)

$$(x^2 + y^2 + z^2 + t^2)^2 = (x^2 + y^2 - z^2 - t^2)^2 + (2xt + 2yz)^2 + (2ty - 2xz)^2,$$

$$(a^2 + b^2 + c^2 + d^2)^2 = (a^2 + b^2 - c^2 - d^2)^2 + (2ad - 2bc)^2 + (2ac + 2bd)^2.$$

These relations can be reduced even farther depending upon what q_1 and q_2 are. A simple observation here is that the simplified relations (16) depend only on the plane containing the square and every other square contained in the same plane has the same simplified relations. This implies that the minimal

square contained in the plane has the sides at most $\sqrt{k_1 k_2}$, where the $k_i^2 = A_i^2 + B_i^2 + C_i^2$ are the primitive versions of (16).

Example 3. We start with two equalities as in (16): $9^2 = 7^2 + 4^2 + 4^2$ and $15^2 = 11^2 + 10^2 + 2^2$. This implies that $45^2 = 33^2 + 30^2 + 6^2 = 35^2 + 20^2 + 20^2$. As in Theorem 2.8, we can take $a = 33$, $a' = 35$, $b = 30$, $b' = 20$, $c = 6$ and $c' = 20$. Then $\Delta_{12} = 1$, $\Delta_{34} = 34$, $\Delta_{13} = 5$, $\Delta_{24} = 25$, $\Delta_{14} = 13$, and $\Delta_{23} = 7$, and the equations of the plane are $34v + 25w + 7t = 0$ and $7u + 5v + w = 0$. The system can be solved easily in terms of u and v : $w = -7u - 5v$ and $t = 25u + 13v$. Then two vectors that generate the lattice in this plane are $(1, 0, -7, 25)$ and $(0, 1, -5, 13)$. Then the ‘‘volume’’ of the fundamental domain is 45. The quadratic form is $Q(u, v) = 675u^2 + 720uv + 195v^2$ or $Q(u, v) = 15(45u^2 + 48uv + 13v^2)$. So, the equation $Q(u, v) = 45\ell$ is equivalent to $(13v + 24u)^2 + 9u^2 = 3(13)\ell$. Then it is clear that v has to be a multiple of 3 and if we take $\ell = 3$, $v = 3$ and $u = -2$ we obtain a solution. This gives $\mathbf{u} = (-2, 3, -1, -11)$, and $v = 6$ and $u = -3$ gives $\mathbf{v} = (-3, 6, -9, 3) = 3(-1, 2, -3, 1)$ (which is indeed a square). This means that the Ehrhart polynomial of this square is $E_{\square}(t) = 3t^2 + 4t + 1 = (t + 1)(3t + 1)$, $t \in \mathbb{N}$. Taking the parameterizations as in Theorem 2.4 for the two equalities, $9^2 = 7^2 + 4^2 + 4^2$ and $15^2 = 11^2 + 10^2 + 2^2$, we observe that this square is covered by the parametrization (15) by taking $a = x = z = 1$, $b = d = t = 2$, $c = 0$, and $y = 3$. It is surprising that such a square with a big size for its side lengths has no lattice points in the interior.

In order to understand better what is happening we will reformulate everything in terms of quaternions.

We remind the reader that the Hamilton quaternion algebra over the real numbers, denoted by $\mathbb{H}(\mathbb{R})$, is the associative unitary algebra given by the requirements:

(I) $\mathbb{H}(\mathbb{R})$ is the free \mathbb{R} -module over the symbols i , j , and k , with 1 the multiplicative unit;

(II) $i^2 = j^2 = k^2 = -1$, $ij = -ji = k$, $jk = -kj = i$ and $ki = -ik = j$.

If $q = x + yi + zj + tk \in \mathbb{H}(\mathbb{R})$ the conjugate of q is $\bar{q} = x - yi - zj - tk$ and the norm of q is $N(q) = x^2 + y^2 + z^2 + t^2$. Some standard notation is then naturally appearing: $Re(q) = x$ and $Im(q) = yi + zj + tk$.

By $\mathbb{H}(\mathbb{Z})$ we denote the subset of quaternions whose components are all integers. We imbed \mathbb{Z}^4 into $\mathbb{H}(\mathbb{Z})$ in the natural way: $(x, y, z, t) \mapsto x + yi + zj + tk$. Also, we will think of \mathbb{R}^3 imbedded in $\mathbb{H}(\mathbb{R})$ in a, more or less, natural way $(y, z, t) \mapsto yi + zj + tk$; in other words, \mathbb{R}^3 is the hyperplane $Re(q) = 0$.

It is known that this norm is multiplicative, *i.e.* $N(q_1 q_2) = N(q_1)N(q_2)$, and $\overline{q_1 q_2} = \overline{q_2} \overline{q_1}$.

For the important results about the arithmetic of $\mathbb{H}(\mathbb{Z})$, we recommend the reader the recent treatment in [5]. Using the same terminology as in [5], a quaternion q is called *odd*, if $N(q)$ is an odd number.

The parametrization in Theorem 2.4 is basically equivalent to

$$(17) \quad n_1i + n_2j + n_3k = q(\epsilon)\bar{q}, \text{ where } q = x + yi + zj + tk, \text{ and } \epsilon \in \{i, j, k\}.$$

In case $\epsilon = k$, $n_1 = 2(xz + yt)$, $n_2 = 2(zt - xy)$, and $n_3 = x^2 - y^2 - z^2 + t^2$.

The parametrization (15) is equivalent to

$$(18)$$

$$u = q_1\epsilon_2\bar{q}_2 \text{ and } v = q_1\epsilon_3\bar{q}_2, \text{ where } q_1 = x + yi + zj + tk, q_2 = a + bi + cj + dk,$$

with ϵ_2 and ϵ_3 fixed in $\{i, j, k\}$. In what follows we are going to take $\epsilon_2 = j$ and $\epsilon_3 = k$. This square is in a plane defined by equations similar to (10) obtained from the two Pythagorean quadruples defined by q_1 and q_2 as in (17) using $\epsilon = \epsilon_1$ such that $\{\epsilon_1, \epsilon_2, \epsilon_3\} = \{i, j, k\}$.

Observation 1. If we substitute $q = \tilde{q}(\alpha + \beta k)$ into (17) for $\epsilon = k$, we see that because $(\alpha + \beta k)k = k(\alpha + \beta k)$, we have

$$\begin{aligned} n_1i + n_2j + n_3k &= \tilde{q}(k)(\alpha + \beta k)(\alpha - \beta k)\tilde{q} = (\alpha^2 + \beta^2)\tilde{q}(k)\tilde{q} \\ &= (\alpha^2 + \beta^2)(\tilde{n}_1i + \tilde{n}_2j + \tilde{n}_3k). \end{aligned}$$

This implies that $n_i = \tilde{n}_i(\alpha^2 + \beta^2)$, $i = 1, 2, 3$, which means that the primitive solutions of the equation (3) have to arise from quaternions q which have no right factors of the form $\alpha + \beta k$. We obtain the same conclusion if $q = \tilde{q}(1 + i)$ or $q = \tilde{q}(1 + j)$, because $(1 + i)k = -j(1 + i)$ and $(1 + j)k = i(1 + j)$. We have the following converse of this observation.

PROPOSITION 2.9. *If in (17), we have $\gcd(n_1, n_2, n_3) = n > 1$, i.e. if $n_1 = 2(xz + yt)$, $n_2 = 2(zt - xy)$ and $n_3 = x^2 - y^2 - z^2 + t^2$, are divisible by n , then $q = x + yi + zj + tk = (x' + y'i + z'j + t'k)\eta$ where $\eta \in \{1 + i, 1 + j, \alpha + k\beta\}$ for some integers α and β .*

Proof. Let us pick a prime p which divides n . Since

$$n_1^2 + n_2^2 = 4[(xz + yt)^2 + (zt - xy)^2] = 4(x^2 + t^2)(y^2 + z^2)$$

is divisible by n^2 it is divisible by p . First, we assume that $p > 2$. Then p divides either $A := x^2 + t^2$ or $B := y^2 + z^2$. Since $n_3 = A - B$ and p divides n_3 , we must have, in fact, A and B both divisible by p . If $p = 2$, then A and B are either both even or both odd. If A and B are both even, then we have the same conclusion as in the case $p > 2$. In the case A and B are both odd, we have, lets say $x = 2x'$, $y = 2y'$, $z = 2z' + 1$, $t = 2t' + 1$. This implies that

$$q = x + yi + zj + tk = 2x' + 2y'i + 2z'j + 2t'k + j + k$$

$$= (x' + y'i + z'j + t'k)(1 - i)(1 + i) + k(1 + i) = \tilde{q}(1 + i)$$

which proves our claim. Similar conclusion can be drawn if x and z , t and y , or t and z are even.

If p is a prime of the form $4k + 3$, then automatically $x = px'$, $z = pz'$, $t = pt'$, and $y = py'$, which shows that $q = x + yi + zj + tk = (x' + y'i + z'j + t'k)p$ and the conclusion of our proposition follows, with $\alpha = p$ and $\beta = 0$.

Finally, if p is a prime of the form $4k + 1$, then $p = \alpha^2 + \beta^2$. Because $A = (x + tk)(x - tk)$ and $\eta = \alpha + \beta k$ is a Gaussian prime integer, it divides $x + tk$ or $x - tk$. Without loss of generality we may assume that η divides $x + tk$. If this is not the case we continue with $\bar{\eta}$. Next we observe that since p divides $xz + yt$ and $zt - xy$, then p divides $(x + tk)(y + zk) = (xy - tz) + (ty + xz)k$. This implies that if $\bar{\eta}$ does not divide $x + tz$, then $\bar{\eta}$ must divide $y + zk$. Since η divides $B = (y + zk)(y - zk)$, then η divides either $y + zk$ or $y - zk$. Let us assume first that η divides $y - zk$. Then we can write

$$q = x + yi + zj + tk = x + tk + i(y - zk) = (x' + t'k)\eta + i(y' + z'k)\eta = \tilde{q}\eta.$$

If η divides $y + zk$, then $\bar{\eta}$ divides $y - zk$. If $\bar{\eta}$ divides $z + tz$ then we proceed as above, with a small change:

$$q = x + yi + zj + tk = x + tk + i(y - zk) = (x' + t'k)\bar{\eta} + i(y' + z'k)\bar{\eta} = \tilde{q}\bar{\eta}.$$

If $\bar{\eta}$ does not divide $x + tz$, then we have shown above that in this case, $\bar{\eta}$ must divide $y + zk$, which is the same thing as η dividing $y - zk$. This puts us in the same position as above. \square

Observation 2. If we substitute $q_1 = \tilde{q}_1(\alpha + \beta i)$ into (18), we see that because $(\alpha + \beta i)k = \alpha k - \beta j$ and $(\alpha + \beta i)j = \alpha j + \beta k$, we have $\tilde{u} = \alpha u - \beta v$ and $\tilde{v} = \beta u + \alpha v$. This is saying that the twin pair \tilde{u} and \tilde{v} is in the same plane as u and v . A similar statement can be obtained if we substitute $q_2 = \tilde{q}_2(\alpha + \beta i)$ into (18).

Let us recall, for the convenience of the reader, the statement of Lemma 2.6.5 in [5], with the only difference that the important factors appear on the right (so, this new statement follows by conjugation and a symmetry type transformation $i \rightarrow -i$, $j \rightarrow -j$, and $k \rightarrow -k$).

LEMMA 2.10 ([5]). *Every integer quaternion $q \in \mathbb{H}(\mathbb{Z})$ has unique factorization*

$$(19) \quad q = 2^{\ell_q} \tilde{q} \eta,$$

where $\tilde{q} \in \mathbb{H}(\mathbb{Z})$ is odd, $\eta_q \in \{1, 1 + i, 1 + j, 1 + k, (1 + j)(1 + i), (1 - k)(1 + i)\}$ and for some non-negative $\ell_q \in \mathbb{Z}$.

This new statement follows by conjugation and a symmetry type transformation $i \rightarrow -i$, $j \rightarrow -j$, and $k \rightarrow -k$ from the original statement in [5].

THEOREM 2.11. (i) *We assume that q_1 and q_2 in the parametrization (18) represented as in (19), are not right-divisible by quaternions of the form $\pi = \alpha + \beta i$, $|\pi| > 1$, then the square in the parametrization (18) is minimal.*

(ii) *The parametrization (18) represents all the integer squares.*

Proof. (i) We observe first that under our hypothesis, $\ell_{q_1} = \ell_{q_2} = 0$ and $\eta_{q_1}, \eta_{q_2} \in \{1 + j, 1 + k\}$ in (19). By way of contradiction, if the construction of the square $\mathcal{S} = \{u, v\}$ in (18) is not minimal, there exists an integer square $\mathcal{S}_0 = \{u_0, v_0\}$ in the same plane in such a way \mathcal{S} can be obtain from \mathcal{S}_0 by a simple transformation which involves two integer parameters α and β : $u = \alpha u_0 + \beta v_0$ and $v = -\beta u_0 + \alpha v_0$. We may assume that $p = \alpha^2 + \beta^2$ is a prime, otherwise we redefine the twin pair \mathcal{S}_0 in such a way this condition is satisfied. Let us first assume that $p > 2$. Since q_1 is not right-divisible by $\eta = \alpha + \beta i$ (which is a prime in $\mathbb{H}(\mathbb{Z})$, η odd quaternion), we conclude that $\gcd(q_1, \bar{\eta})_r = 1$ (the right greatest-common divisor as defined in [5]). By Theorem 2.6.6 in [5], which is the equivalent of Bézout's relation in $\mathbb{H}(\mathbb{Z})$, we can find γ and δ in $\mathbb{H}(\mathbb{Z}[\frac{1}{2}])$ such that

$$\gamma q_1 + \delta \bar{\eta} = 1.$$

If $p = 2$, then $|u|^2 = |v|^2 = N(q_1)N(q_2) = 2|u_0|^2$ and so 2 divides $N(q_1)$ or $N(q_2)$. Using Lemma 2.10 we conclude that q_1 is of the form $q'_1(1 + k)$ or $q'_1(1 + j)$ with q'_1 odd. So we can still obtain a Bézout's relation as above, with q'_1 and $\bar{\eta}$, which can be easily transformed into one for q_1 and $\bar{\eta}$.

Solving for u_0 , the above system, in terms of u and v , we have

$$u_0 = (1/p)(\alpha u - \beta v) = (1/p)q_1(\alpha + i\beta)k\bar{q}_2.$$

Multiplying on the right with $(1/p)(\alpha + i\beta)k\bar{q}_2$ the above Bézout's relation, we obtain

$$\gamma u_0 + \delta k\bar{q}_2 = (1/p)(\alpha + i\beta)k\bar{q}_2.$$

From this relation we see that $q_2 = \tilde{q}_2(\alpha - i\beta)$ for some $\tilde{q}_2 \in \mathbb{H}(\mathbb{Z}[\frac{1}{2}])$. Hence $2^k q_2 = q'_2(\alpha - i\beta)$ for some $q'_2 \in \mathbb{H}(\mathbb{Z})$. Using Lemma 2.10 again, we conclude that q_2 is right divisible by $\alpha - i\beta$ which is in contradiction with our hypothesis. Therefore, it must be true that the $\mathcal{S} = \{u, v\}$ in (18) is minimal.

(ii) Let us start with an arbitrary square, \mathcal{S}_0 . We then use Theorem 2.7 to find the reduced equations of the plane that it is contained in, as those given by (10). Then we can apply the Theorem 2.4 twice, and finally use the two sets of parameters (8 in total) to construct a square \mathcal{S}_1 , with formulae (15). The quaternions involved have the property in part (i) (both odd), and so the square \mathcal{S}_1 is minimal. Then \mathcal{S}_0 can be written in terms of \mathcal{S}_1 using two

numbers α and β . Using the substitutions described before the statement of the theorem, we can write \mathcal{S}_0 as in (18) by multiplying either of the quaternions by $\alpha + \beta i$. \square

THEOREM 2.12. *Assuming we have two odd integer quaternions q_1 and q_2 as in (ii) in Theorem 2.11, then the fundamental domain of the integer lattice containing the square $\{u, v\}$ in (18), has “volume” equal to $V = \text{lcm}(N(q_1), N(q_2))$. The Ehrhart polynomial of the square (18) is*

$$E_{\square}(t) = \text{gcd}(N(q_1), N(q_2))t^2 + (D_1 + D_2)t + 1,$$

where $D_1 = \text{gcd}(u_1, u_2, u_3, u_4)$ and $D_2 = \text{gcd}(v_1, v_2, v_3, v_4)$.

Proof. This follows from Theorem 2.7 (part (ii)), Proposition 2.9 and the observation that $\text{lcm}(N(q_1), N(q_2))^2 = a^2 + b^2 + c^2 = a'^2 + b'^2 + c'^2$ for some integers a, b, c, a', b', c' with $\text{gcd}(a, b, c, a', b', c') = 1$. \square

Observation 3. To get back in dimension 3, we identified \mathbb{R}^3 with the subspace of quaternions q for which $\text{Re}(q) = 0$. We have seen that only one Pythagorean quadruple equation is necessary: $\Delta_{13} = \Delta_{12} = 0$ which implies $\Delta_{14} = 0$. To accomplish this, we can take $q_1 = q_2 = q$ where q is odd. This implies that the most general square in dimension 3 is of the form (4) combined with the usual variations from dimension 2 in the respective plane: $\tilde{u} = \alpha u - \beta v$ and $\tilde{v} = \beta u + \alpha v$ with $\{u, v\}$ as in (4).

There seems to be plenty of numerical evidence that the sequence of almost perfect squares in dimension four is the set of all non-negative integers. Let us denote the set of all such numbers by \mathcal{APS}_4 . We include some partial result in this direction.

THEOREM 2.13. (i) *The set \mathcal{APS}_4 contains all odd numbers.*

(ii) *The set \mathcal{APS}_4 contains all even numbers of the form $p - 1$ with p a prime $p \geq 11$.*

Proof. (i) Let us take an odd k , $k \geq 1$, and a primitive Pythagorean quadruple representation $k^2 = a^2 + b^2 + c^2$. Without loss of generality we may assume that a is odd and both b and c are even. If we multiply this equality by 2, we obtain $2k^2 = 2a^2 + 2b^2 + 2c^2 = 2c^2 + (a+b)^2 + (a-b)^2$. Hence we can build the square $u = (k, k, 0, 0)$ and $v = (c, -c, a+b, a-b)$. It is clear that $D_1 = k$ and $D_2 = 1$ ($a-b$ and $a+b$ are odd) as defined in Theorem 2.12. Since the sides have length $k\sqrt{2}$ and the two equations defining the plane containing $\{u, v\}$ are $k^2 = (-c)^2 + b^2 + a^2 = (-c)^2 + a^2 + b^2$, we conclude by Theorem 2.12 that the Ehrhart polynomial is $E_{u,v}(t) = 2kt^2 + (k+1)t + 1$. Hence $E_{u,v}(-1) = k$. This proves the first statement.

(ii) If p is a prime $p \geq 11$, then we can find two distinct primitive Pythagorean quadruple representations $p^2 = a^2 + b^2 + c^2 = a'^2 + b'^2 + c'^2$ and construct a square as in (18), whose Ehrhart polynomial is $E(t) = pt^2 + (D_1 + D_2)t + 1$, where D_i divide p . It is not possible to have $D_i = p$ since this puts us in the position where one of the vectors defining the square, say u , is of the form $u = (p, 0, 0, 0)$. This leads essentially to only one equation which defines the plane containing $\{u, v\}$, but we have two different equations that define the plane. Hence, since $E(-1) = p - 1$ the claim in (ii) follows. \square

Numerical evidence shows that given an odd $k \geq 9$, one can find a square whose Ehrhart polynomial is $E(t) = kt^2 + 2t + 1$. For instance, for $k = 2015$ we found $u = (-836, 584, -1592, -697)$ and $v = (-506, 1414, 203, 1328)$ that does the job.

3. CUBES

If a general lattice cube in \mathbb{R}^3 is given by the orthogonal matrix

$$(20) \quad C_\ell = \frac{1}{\ell} \begin{bmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{bmatrix},$$

with a_i, b_i and c_i integers satisfying $a_i a_j + b_i b_j + c_i c_j = \delta_{i,j} \ell^2$ for all i, j in $\{1, 2, 3\}$, we define $d_i := \gcd(a_i, b_i, c_i)$. It is clear that the d_i are divisors of ℓ . Assuming that the matrix in (20) is irreducible (no smaller ℓ can be used, which implies ℓ odd), then we have shown in [19] the following.

THEOREM 3.1. *Given a cube C_ℓ constructed from a matrix as in (20), its Ehrhart polynomial is given by*

$$(21) \quad L(C_\ell, t) = \begin{aligned} & \ell^3 t^3 + \ell(d_1 + d_2 + d_3)t^2 + (d_1 + d_2 + d_3)t + 1 \text{ or} \\ & (\ell t + 1)[\ell^2 t^2 + (d_1 + d_2 + d_3 - \ell)t + 1] \end{aligned}, \quad t \in \mathbb{N}.$$

Similar formula takes place for the cubes in dimension four. In general, a square can be completed in various ways to a cube and even to a hypercube. Let us assume that the cube is given by the first three vectors in the rows of the following matrix

$$(22) \quad \begin{bmatrix} a_1 & b_1 & c_1 & d_1 \\ a_2 & b_2 & c_2 & d_2 \\ a_3 & b_3 & c_3 & d_3 \\ a_4 & b_4 & c_4 & d_4 \end{bmatrix},$$

with a_i, b_i, c_i and d_i integers $a_i a_j + b_i b_j + c_i c_j + d_i d_j = \delta_{i,j} \ell$ for all i, j in $\{1, 2, 3, 4\}$. In this case, we define $D_i := \gcd(a_i, b_i, c_i, d_i)$ and ζ_{ij} be the greatest common divisor of all the 2-by-2 determinants of the matrix

$$\begin{bmatrix} a_i & b_i & c_i & d_i \\ a_j & b_j & c_j & d_j \end{bmatrix},$$

for every i and j with $1 \leq i < j \leq 3$.

THEOREM 3.2. *Given a cube C_ℓ constructed from a matrix as in (22), its Ehrhart polynomial is given by*

$$(23) \quad L(C_\ell, t) = \ell D_4 t^3 + \Delta t^2 + \Delta' t + 1, \quad t \in \mathbb{N},$$

where $\Delta := \zeta_{12} + \zeta_{13} + \zeta_{23}$ and $\Delta' := D_1 + D_2 + D_3$.

Proof. For the first coefficient the volume of the cube is $\ell\sqrt{\ell}$ and the volume of the fundamental domain of the lattice containing it is equal to $\sqrt{a_4^2 + b_4^2 + c_4^2 + d_4^2}/D_4 = \frac{\sqrt{\ell}}{D_4}$. For the second coefficient, the area of each of the six faces is ℓ . Using the Theorem 2.11 to find the “volume” of the fundamental domain of the lattice containing the faces determined by rows i and j , we obtain ℓ/δ_{ij} , which implies the stated value of Δ . Finally, for the coefficient Δ' , we proceed like in the proof of Theorem 3.1 [19]. The main idea is essentially based on the fact that the topological equivalent in our cube to $[0, 1]^3$ (containing κ lattice points) has the fundamental domain property: the dilation with a factor t contains exactly $t^3 \kappa$ lattice points. This implies that $\kappa = \ell D_4$ and so

$$\ell D_4 = 1 + (\ell D_4 - \Delta + \Delta' - 1) + \sum_{ij} (\zeta_{ij} - (D_i + D_j) + 1) + \sum_i (D_i - 1).$$

Solving for Δ' we obtain the stated value in the theorem. \square

3.1. Hypercubes

Hypercubes in dimension four can be constructed using quaternion techniques described earlier. A few orthogonal matrices in dimension 4, together with the Ehrhart polynomial associated to the corresponding hypercube, are included next:

$$\frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 & 0 & 0 \\ 1 & -10 & 0 & \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & -1 \end{bmatrix}, \quad \frac{1}{\sqrt{3}} \begin{bmatrix} 1 & 1 & 1 & 0 \\ -1 & 1 & 0 & 1 \\ 0 & -11 & 1 & \\ -1 & 0 & 1 & -1 \end{bmatrix}, \quad \frac{1}{2} \begin{bmatrix} 1 & 1 & 1 & -1 \\ -1 & 1 & 1 & 1 \\ 1 & -1 & 1 & 1 \\ 1 & 1 & -1 & 1 \end{bmatrix},$$

$$4t^4 + 8t^3 + 8t^2 + 4t + 1 = (2t^2 + 2t + 1)^2, \quad 9t^4 + 12t^3 + 6t^2 + 4t + 1 = (t + 1)(3t + 1)(3t^2 + 1), \quad 16t^4 + 16t^3 + 12t^2 + 4t + 1 = (1 + 2t + 4t^2)^2$$

$$\frac{1}{\sqrt{5}} \begin{bmatrix} 2 & 1 & 0 & 0 \\ 1 & -20 & 0 & \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 1 & -2 \end{bmatrix}, \quad \frac{1}{\sqrt{6}} \begin{bmatrix} 2 & 1 & 1 & 0 \\ 1 & -2 & 0 & 1 \\ 1 & 0 & -2 & -1 \\ 0 & 1 & -1 & 2 \end{bmatrix}, \quad \frac{1}{\sqrt{7}} \begin{bmatrix} 2 & 1 & 1 & 1 \\ 1 & -2 & -1 & 1 \\ 1 & 1 & -2 & -1 \\ 1 & -1 & 1 & -2 \end{bmatrix},$$

$$25t^4 + 20t^3 + 14t^2 + 4t + 1 \quad 36t^4 + 24t^3 + 8t^2 + 4t + 1 \quad 49t^4 + 28t^3 + 6t^2 + 4t + 1$$

$$= (1 + 2t + 5t^2)^2$$

$$\frac{1}{3} \begin{bmatrix} 30 & 0 & 0 \\ 02 & 2 & 1 \\ 02 & -1 & -2 \\ 01 & -2 & 2 \end{bmatrix}, \quad \frac{1}{3} \begin{bmatrix} 2 & 2 & 1 & 0 \\ 2 & -2 & 0 & 1 \\ 1 & 0 & -2 & -2 \\ 0 & 1 & -2 & 2 \end{bmatrix}, \quad \frac{1}{\sqrt{10}} \begin{bmatrix} 2 & 2 & 1 & 1 \\ 2 & -2 & -1 & 1 \\ 1 & 1 & -2 & -2 \\ 1 & -1 & 2 & -2 \end{bmatrix},$$

$$81t^4 + 54t^3 + 18t^2 + 6t + 1 \quad 81t^4 + 36t^3 + 6t^2 + 4t + 1 \quad 100t^4 + 40t^3 + 16t^2 + 4t + 1$$

$$= (3t + 1)^2(9t^2 + 1)$$

$$\frac{1}{\sqrt{10}} \begin{bmatrix} 3 & -10 & 0 \\ 1 & 3 & 0 & 0 \\ 0 & 0 & 3 & 1 \\ 0 & 0 & 1 & -3 \end{bmatrix}, \quad \frac{1}{\sqrt{11}} \begin{bmatrix} 3 & 1 & 1 & 0 \\ 1 & -3 & 0 & 1 \\ 1 & 0 & -3 & -1 \\ 0 & 1 & -1 & 3 \end{bmatrix}, \quad \text{and} \quad \frac{1}{\sqrt{13}} \begin{bmatrix} 2 & 2 & 2 & 1 \\ 2 & -2 & 1 & -2 \\ 2 & -1 & -2 & 2 \\ 1 & 2 & -2 & -2 \end{bmatrix},$$

$$100t^4 + 40t^3 + 24t^2 + 6t + 1 \quad 121t^4 + 44t^3 + 6t^2 + 4t + 1 \quad 169t^4 + 53t^3 + 6t^2 + 4t + 1$$

$$= (10t^2 + 2t + 1)^2$$

We see that several of these examples are cross polytopes (squares in two dimensions) and indeed the resulting polynomial is the product of the smaller degree polynomials involved. Let us assume that in general we have

$$(24) \quad H_\ell = \frac{1}{\sqrt{\ell}} \begin{bmatrix} a_1 b_1 c_1 d_1 \\ a_2 b_2 c_2 d_2 \\ a_3 b_3 c_3 d_3 \\ a_4 b_4 c_4 d_4 \end{bmatrix},$$

where $a_i a_j + b_i b_j + c_i c_j + d_i d_j = \delta_{ij} \ell$ for $i, j \in \{1, 2, 3, 4\}$. We define $D_i = \gcd(a_i, b_i, c_i, d_i)$ for $i \in \{1, 2, 3, 4\}$ and assume that H_ℓ is irreducible, *i.e.*, $\gcd(D_1, D_2, D_3, D_4) = 1$. The Ehrhart polynomial associated to the hypercube constructed in a natural way using the orthogonal matrix H_ℓ is then

$$(25) \quad E_{H(\ell)}(t) = \ell^2 t^4 + \alpha_1 t^3 + \alpha_2 t^2 + \alpha_3 t + 1.$$

PROPOSITION 3.3. *With the notation introduced earlier $\alpha_1 = \ell(D_1 + D_2 + D_3 + D_4)$.*

Proof. Let us look at a 3-dimensional face, say generated by first three rows of (22). Its volume is $\ell\sqrt{\ell}$ and the volume of the lattice containing it is equal to $\sqrt{a_4^2 + b_4^2 + c_4^2 + d_4^2}/D_4 = \frac{\sqrt{\ell}}{D_4}$. Since there are eight such faces the general theory gives the value claimed. \square

THEOREM 3.4. *We have*

$$(26) \quad \alpha_2 = \alpha_3 + \delta - \Delta,$$

where $\delta = \sum_{1 \leq i < j \leq 4} \zeta_{ij}$ and $\Delta = D_1 + D_2 + D_3 + D_4$.

Proof. We are going to use the same property of the part of hypercube that is topologically equivalent to $[0, 1]^4$, as in the proof of Theorem 3.2. The balancing of points in each of the corresponding sets of the partition

$$[0, 1]^4 \\ = (0, 0, 0, 0) \cup (0, 1)^4 \quad \bigcup_{\text{all four 3D cubes}} (0, 1)^3 \quad \bigcup_{\text{all six 2D faces}} (0, 1)^2 \quad \bigcup_{\text{all four 1D sides}} (0, 1)$$

gives (using Theorem 3.2)

$$\ell^2 = 1 + (\ell^2 - \alpha_1 + \alpha_2 - \alpha_3 + 1) + (\ell\Delta - 2\delta + 4\Delta - 4) + \sum_{1 \leq i < j \leq 4} (\zeta_{ij} - (D_i + D_j) + 1) + \Delta - 4.$$

From this, one can easily derive (26). \square

Looking at the examples we have so far, we notice that $\alpha_3 = \Delta$, and so the Ehrhart polynomial of an hypercube would result into the following simple form

$$(27) \quad E_{H(\ell)}(t) = \ell^2 t^4 + \ell \Delta t^3 + \delta t^2 + \Delta t + 1.$$

This is indeed the case since it follows from Theorem 9.9 in [2]. Of course, one may want to look into all these concepts for dimensions bigger than four, although it seems that very little is known about the essentially new related objects in dimensions five or beyond. For other further developments, it may be useful to analyze the roots or calculate better bounds for them, such as those in [3], for the type of polytopes studied here.

REFERENCES

- [1] A. Barvinok, *Integer Points in Polyhedra*. Zur. Lect. Adv. Math., European Mathematical Society, Zürich, 2008.
- [2] M. Beck and S. Robins, *Computing the Continuous Discretely: Integer-Point Enumeration in Polyhedra*. Undergrad. Texts Math., Springer-Verlag, New York, 2015.
- [3] M. Beck, J.A. DeLoera, M. Develin, J. Pfeifle and R. Stanley, *Coefficients and roots of Ehrhart polynomials*. Contemp. Math. **374** (2005), 15–36.
- [4] S. Cooper and M. Hirschhorn, *On the number of primitive representations of integers as a sum of squares*. Ramanujan J. **13** (2007), 7–25.
- [5] G. Davidoff, P. Sarnak and A. Valente, *Elementary Number Theory, Group Theory and Ramanujan Graphs*. London Math. Soc. Stud. Texts **55**, Cambridge, 2003.
- [6] M. Beck, R. Diaz and S. Robins, *The Frobenius problem, rational polytopes, and Fourier-Dedekind sums*. J. Number Theory **96** (2002), 1–21.

- [7] R. Diaz and S. Robins, *The Ehrhart polynomial of a lattice polytope*. Ann. of Math. **145** (1997), 503–518.
- [8] E. Ehrhart, *Sur les polygones et les polyèdres réguliers entiers*. Enseign. Math. (2) **5** (1959), 81–85.
- [9] E. Ehrhart, *Sur les polyèdres rationnels homothétiques à n dimensions*. C.R. Acad. Sci. Paris **154** (1962), 616–618.
- [10] E. Ehrhart, *Sur un problème de géométrie diophantienne linéaire. I. Polyèdres et réseaux*. J. Reine Angew. Math. **226** (1967), 1–29.
- [11] E. Ehrhart, *Solution to problem 6179 [1977, 744]*. Amer. Math. Monthly (1980), 826–827.
- [12] S. Frischa and L. Vasersteinba, *Parametrization of Pythagorean triples by a single triple of polynomials*. J. Pure Appl. Algebra **212** (2008), 271–274.
- [13] L.M. Goswick, E.W. Kiss, G. Moussong and N. Simányi, *Sums of squares and orthogonal integral vectors*. J. Number Theory **132** (2012), 37–53.
- [14] R.L. Graham, J.C. Lagarias, C.L. Mallows, A.R. Wilks and C.H. Yan, *Apollonian circle packings: number theory*. J. Number Theory **100** (2003), 1–45.
- [15] W. Hürlimann, *Exact and asymptotic evaluation of the number of distinct primitive cuboids*. J. Integer Seq. **18** (2015), 2, Article 15.2.5
- [16] W. Hürlimann, *On the number of primitive of Pythagorean quintuples*. Journal of Algebra, Number Theory: Advances and Applications **13** (2015), 1, 13–28.
- [17] E.J. Ionascu, *Equilateral triangles in \mathbb{Z}^4* . Vietnam J. Math. **43** (2015), 3, 525–539.
- [18] E.J. Ionascu, *New parametrization of $A^2 + B^2 + C^2 = 3D^2$ and Lagrange's four-square theorem*. An. Ştiinţ. Univ. Al. I. Cuza Iaşi. Mat. (N.S.) **62** (2016), 3, 823–833.
- [19] E.J. Ionascu, *Lattice platonic solids and their Ehrhart polynomial*. Acta Math. Univ. Comenian. **82** (2013), 1, 147–158.
- [20] E.W. Kiss and P. Kutas, *Cubes of integral vectors in dimension four. (English summary)*. Studia Sci. Math. Hungar. **49** (2012), 4, 525–537.
- [21] A. Osborne and H. Liebeck, *Orthogonal bases of \mathbb{R}^3 with integer coordinates and integer lengths*. Amer. Math. Monthly **96** (1989), 49–53.
- [22] A. Osborne and H. Liebeck, *The generation of all rational orthogonal matrices*. Amer. Math. Monthly **98** (1991), 131–133.
- [23] S. Sam, *A bijective proof for a theorem of Ehrhart*. Amer. Math. Monthly **116** (2009), 8, 688–701.
- [24] I.J. Schoenberg, *Regular simplices and quadratic forms*. J. Lond. Math. Soc. **12** (1937), 48–55.
- [25] R. Spira, *The Diophantine equation $x^2 + y^2 + z^2 = m^2$* . Amer. Math. Monthly **69** (1962), 360–65.

Received 1 December 2017

Columbus State University
 Department of Mathematics
 4225 University Avenue
 Columbus, GA 31907
 Honarific Member
 of the Romanian Institute
 of Mathematics “Simion Stoilow”
 ionascu@columbusstate.edu
 math@ejionascu.ro