

REFLECTIONS ON TRISECTION GENUS

MICHELLE CHU and STEPHAN TILLMANN

The Heegaard genus of a 3-manifold, as well as the growth of Heegaard genus in its finite sheeted covering spaces, has extensively been studied in terms of algebraic, geometric and topological properties of the 3-manifold. This note shows that analogous results concerning the trisection genus of a smooth, orientable 4-manifold have more general answers than their counterparts for 3-manifolds. In the case of hyperbolic 4-manifolds, upper and lower bounds are given in terms of volume and a trisection of the Davis manifold is described.

AMS 2010 Subject Classification: 57Q15, 57N13, 51M10, 57M50.

Key words: trisection, trisection genus, stable trisection genus, triangulation complexity, rank of group, Davis manifold.

1. BOUNDS ON TRISECTION GENUS

Gay and Kirby's construction of a *trisection* for arbitrary smooth, oriented closed 4-manifolds [6] defines a decomposition of the 4-manifold into three 4-dimensional 1-handlebodies glued along their boundaries in the following way: Each handlebody is a boundary connected sum of copies of $S^1 \times B^3$, and has boundary a connected sum of copies of $S^1 \times S^2$ (here, B^i denotes the i -dimensional ball and S^j denotes the j -dimensional sphere). The triple intersection of the 4-dimensional 1-handlebodies is a closed orientable surface Σ , called the *central surface*, which divides each of their boundaries into two 3-dimensional 1-handlebodies (and hence is a Heegaard surface). These 3-dimensional 1-handlebodies are precisely the intersections of pairs of the 4-dimensional 1-handlebodies.

A trisection naturally gives rise to a quadruple of non-negative integers $(g; g_0, g_1, g_2)$, encoding the genus g of the central surface Σ and the genera g_0 , g_1 , and g_2 of the three 4-dimensional 1-handlebodies. We allow the genera to be distinct following [14, 19]. The *trisection genus* of M , denoted $g(M)$, is the minimal genus of a central surface in any trisection of M . A trisection with $g(\Sigma) = g(M)$ is called a *minimal genus trisection*.

If M has a $(g; g_0, g_1, g_2)$ -trisection, then $\chi(M) = 2 + g - g_0 - g_1 - g_2$. Since the fundamental group of each 4-dimensional 1-handlebody surjects onto the fundamental group of M , we have $g_k \geq \beta_1(M)$. Combining the above equation with $\chi(M) = 2 - 2\beta_1(M) + \beta_2(M)$ therefore gives

$$g(M) \geq \beta_1(M) + \beta_2(M).$$

This is an equality for the six manifolds with trisection genus at most two [15], and for all standard simply connected 4-manifolds [20]. Castro and Ozbagci [4] constructed trisections of genus $2g+2$ of the twisted bundles $S_g \tilde{\times} S^2$, where S_g is a closed, connected, oriented surface of genus g . By the above inequality, these are all minimal and so $g(S_g \tilde{\times} S^2) = 2g + 2$. More generally, using thin-position arguments, Aranda [2] constructs trisections of all orientable S^2 -bundles over closed (orientable or non-orientable) surfaces. All of these are minimal due to the above bound (using \mathbb{Z}_2 -coefficients in the case of non-orientable base). Meier [13] produces infinitely many 4-manifolds admitting a minimal genus $(g; k, k, k)$ -trisection for each $g \geq 3$ and each $0 < k < g - 1$. Minimality is shown in [13] using observations that are equivalent to the ones given in this paper.

We have $g_k \geq \text{rk } \pi_1(M) \geq \beta_1(M)$, where $\text{rk}(\pi_1(M))$ is the minimal number of generators of the fundamental group of M . Hence, lower bounds on the rank give lower bounds on trisection genus, and, conversely, upper bounds on the trisection genus give upper bounds on the rank. Moreover, $g(M) \geq \chi(M) - 2 + 3 \text{rk}(\pi_1(M))$.

Question 1. Given any finitely presented group G , is there a smooth, oriented closed 4-manifold M with $\pi_1(M) = G$ such that $g(M) = \chi(M) - 2 + 3 \text{rk}(G)$?

The question has a positive answer for surface groups (using the above examples of $S_g \tilde{\times} S^2$) and the infinite cyclic group (by virtue of $S^1 \times S^3$). The construction of trisections for spun 4-manifolds by Meier [13] shows that the above question also has a positive answer for every group that is the fundamental group of a 3-manifold satisfying the rank versus genus conjecture. It thus has a positive answer for free products of these groups (in particular, finitely generated free groups), by taking connected sums of these manifolds and application of Grushko's theorem [8, 17].

We now turn to an upper bound. Let $\sigma(M)$ be the minimal number of 4-simplices in a (possibly singular) triangulation of M . We call $\sigma(M)$ the *triangulation complexity* of M . Then the upper bound on trisection genus recently given in [3] using triangulations gives:

$$60 \sigma(M) \geq g(M).$$

This note gives some simple consequences of these upper and lower bounds. Our starting point is the following, which at once gives upper and lower bounds on trisection genus, *and* a lower bound on the triangulation complexity of a 4-manifold.

PROPOSITION 2. *Let M be a smooth orientable 4-manifold and assume that M is not diffeomorphic to S^4 . Then*

$$60\sigma(M) \geq g(M) \geq \frac{1}{3}|\chi(M)|.$$

The proof is simple. We have $\chi(M) = 2 + g - g_0 - g_1 - g_2$. Hence $g(M) \geq \chi(M) - 2$. In particular, the lower bound holds when $\chi(M) \geq 3$. If $\chi(M) > 0$ and $\beta_1(M) > 0$, then $g(M) \geq \chi(M)$. Now suppose $1 \leq \chi(M) \leq 2$, $\beta_1(M) = 0$ and $M \neq S^4$. The classification of trisections of genus 2 due to Meier and Zupan [15], then implies $g(M) \geq 3 \geq \chi(M)$. So the lower bound holds in the case of positive Euler characteristic. Moreover, if $\mathbb{C}P^2$ is excluded from the statement, one may replace $1/3$ by $1/2$.

Combining $\chi(M) = 2 - 2\beta_1(M) + \beta_2(M)$ with the bound $g(M) \geq \beta_1(M)$ gives $g(M) \geq 1 - \frac{1}{2}\chi(M)$. So if $\chi(M) \leq 0$, then $g(M) \geq \frac{1}{2}|\chi(M)| + 1$. This is sharp for $S^1 \times S^3$, where $\chi(S^1 \times S^3) = 0$ and $g(S^1 \times S^3) = 1$. \square

1.1. Stable trisection genus

We now turn to finite sheeted covers. In particular, we are interested in the case where there are finite covers of arbitrarily large degree, but our results also apply when there are only finitely many such covers. If M is hyperbolic, then residual finiteness implies that there are finite covers of arbitrarily large degree. If $\chi(M) < 0$, then $\beta_1(M) > 0$, and hence there are finite cyclic covers of any degree. Since the only manifold covered by S^4 is the non-orientable manifold $\mathbb{R}P^4$ [9, Section 12.1], we have:

COROLLARY 3. *Let M be a smooth orientable 4-manifold and assume that M is not diffeomorphic to S^4 . Suppose $N \rightarrow M$ is a finite sheeted cover of degree d . Then $g(N)$ is $O(d)$. Specifically,*

$$60\sigma(M) d \geq g(N) \geq \frac{1}{3}|\chi(M)| d.$$

Moreover, if $\chi(M) \neq 0$, then $g(N)$ is $\Theta(d)$.

Another immediate corollary is in terms of the characteristic functions of Milnor and Thurston [16]. Denote

$$\sigma_\infty(M) = \inf \left\{ \frac{\sigma(N)}{d} \mid N \rightarrow M \text{ a degree } d \text{ cover} \right\}$$

and define the *stable trisection genus* by

$$g_\infty(M) = \inf \left\{ \frac{g(N)}{d} \mid N \rightarrow M \text{ a degree } d \text{ cover} \right\}.$$

For instance $g_\infty(S^1 \times S^3) = 0$, $g_\infty(T^4) = 0$, $g_\infty(S_g \times S^2) = 2g - 2 = -\chi(S_g) = g_\infty(S_g \tilde{\times} S^2)$ if $g \geq 1$.

The definition of stable trisection genus and Proposition 2 imply:

COROLLARY 4. *Let M be a smooth orientable 4-manifold and assume that M is not diffeomorphic to S^4 . Then*

$$60 \sigma_\infty(M) \geq g_\infty(M) \geq \frac{1}{3} |\chi(M)|.$$

Betti numbers grow at most linearly in the degree, since the homology is carried by a lifted triangulation. Moreover, the equality $\chi(M) = 2 - 2\beta_1(M) + \beta_2(M)$ implies that $\beta_2(M)$ must grow linearly if $\chi(M) > 0$, and $\beta_1(M)$ must grow linearly if $\chi(M) < 0$. The question of whether the growth is linear or sublinear for the other Betti number is governed by the L^2 -Betti number $b_k^{(2)}(\tilde{M}; \pi_1(M))$ due to work of Lück [12]. Sublinear growth of β_1 and linear growth of β_2 occurs in covers of hyperbolic 4-manifolds, due to positive Euler characteristic and the work of Lück [12]. Linear growth of both β_1 and β_2 occurs in cyclic covers of $(S_g \times S^2) \# (S^1 \times S^3)$, where one unwraps the S^1 -factor. Linear growth of β_1 and sublinear growth of β_2 occurs in covers of $S_g \times S^2$, where β_2 has constant value 2.

1.2. Geometric consequences

In dimension four, Euler characteristic is related to signature (in case the manifold is Einstein), volume or injectivity radius (in case the manifold is hyperbolic). This gives simple applications of the previous results. For instance, the Gromov-Hitchin-Thorpe inequality [7, 10] implies

COROLLARY 5. *Let M be a closed Einstein 4-manifold not diffeomorphic to S^4 . Then*

$$g(M) \geq \frac{1}{2} |\text{sign}(M)| + \frac{1}{7776\pi^2} \|M\|,$$

where $\text{sign}(M)$ is the signature of the intersection form and $\|M\|$ is the Gromov norm.

For congruence covers of arithmetic hyperbolic 3-manifolds, Lackenby [11, Corollary 1.6] bounds Heegaard genus in terms of volume. His lower bound is established in terms of the Cheeger constant, which is uniformly bounded below

for congruence covers. The following counterpart for trisection genus is both more general and has an elementary proof. The following two observations give some answers to Problem 1.24 in [1].

COROLLARY 6. *Let M be a closed hyperbolic 4-manifold. Then there is a positive constant $C = C(M)$ such that for any finite cover $M_i \rightarrow M$,*

$$C \operatorname{Volume}(M_i) \geq g(M_i) \geq \frac{3}{8\pi^2} \operatorname{Volume}(M_i).$$

Proof. As noted in the proof of Proposition 2, $g(M_i) \geq \chi(M_i) - 2$. Since $\chi(M_i)$ is positive and even and $g(M_i) \geq 3$, in fact $g(M_i) \geq \frac{1}{2}\chi(M_i)$. This together with $\operatorname{Volume}(M_i) = \frac{4}{3}\pi^2\chi(M_i)$ gives the lower bound. The upper bound follows directly from Corollary 3 by taking $C(M) = 60 \frac{\sigma(M)}{\operatorname{Volume}(M)}$. \square

Since volume bounds injectivity radius, we have:

COROLLARY 7. *Let M be a closed hyperbolic 4-manifold and let $\operatorname{inj}(M)$ denote its injectivity radius. Then there is a universal positive constant C' such that $g(M) \geq C' \cdot \operatorname{inj}(M)$.*

1.3. The Davis manifold

We conclude this paper by describing trisections of the Davis manifold [5]. The Davis manifold M_D is obtained by identifying opposite pairs of faces of the 120-cell with dihedral angles $\frac{2\pi}{5}$ in hyperbolic space. The boundary of the 120-cell has a cell decomposition with 120 dodecahedra, 720 pentagons, 1200 edges and 600 vertices. The face identifications leave us in M_D with 60 dodecahedra, 144 pentagons, 60 edges and 1 vertex. A natural triangulation of M_D is obtained as follows. Place a vertex v_4 at the centre of the 120-cell, a vertex v_3 at the centre of a dodecahedral face, a vertex v_2 at the barycentre of a pentagonal face thereof, a vertex v_1 at the barycentre of an edge of this, and a vertex v_0 at a vertex of this edge. This gives a Coxeter 4-simplex, usually denoted Δ_3 , and the 120-cell is tiled by $(120)^2 = 14,400$ 4-simplices that are copies of Δ_3 . In particular, there is a group Γ of order $(120)^2$ acting on M_D with M_D/Γ equal to the simplex orbifold with underlying space Δ_3 . Ratcliffe and Tschantz [18] computed $\beta_1(M_D) = 24$ and $\beta_2(M_D) = 72$. Whence

$$864,000 = 60 \cdot (120)^2 \geq g(M_D) \geq 96,$$

using the a priori bounds. We now describe how in situations such as the Coxeter construction of the Davis manifold, the upper bound can be greatly improved using the techniques of [3, 19].

The idea is to partition the vertices of the triangulation into three sets, such that each 4-simplex meets one of them in one vertex and each of the other two sets in two vertices. This partition is then used to define a piece-wise linear map to the 4-simplex. It is shown in [3, 19] that applying certain 2–4 bistellar moves to such a tricoloured triangulation gives a new triangulation with the property that the pull-back of the natural cubulation of the 2–simplex defines a trisection of the 4-manifold. We describe this for the case of the Davis manifold, and refer the reader to the treatment in [3] for more details.

Define a partition of the vertices of the 4-simplices by $S_0 = \Gamma \cdot v_0 \cup \Gamma \cdot v_1$, $S_1 = \Gamma \cdot v_2$ and $S_2 = \Gamma \cdot v_3 \cup \Gamma \cdot v_4$. Then the graphs Γ_k spanned by S_k in the 1-skeleton of the triangulation of M_D have the following properties. The graph Γ_1 is the quotient of the 1-skeleton of the 120-cell and hence a bouquet of 60 circles. The graph Γ_2 is the dual 1-skeleton of the cellulation of M_D arising from the 120-cell and hence also a bouquet of 60 circles. The graph Γ_2 consists of 144 isolated vertices. Each 4-simplex has two vertices in each S_0 and S_2 , and one in S_1 . Hence it meets a unique 4-simplex in the tetrahedral face with all vertices in $S_0 \cup S_1$. This gives a decomposition of M_D in double-4-simplices. As described in [3, Construction 3], we now apply 2–4 bistellar moves to each of these double 4-simplices. This increases the number of pentachora to 28,800. The graphs Γ_0 and Γ_1 are not changed, and Γ_2 turns into a connected graph with 144 vertices and 7,200 edges. Hence Γ_2 is homotopic to a bouquet of 7,057 circles.

We now compute the Euler characteristic of the central surface Σ . We obtain one square for each pentachoron, hence there are 28,800 squares. The number of vertices of the surface equals the number of triangles in the triangulation that have vertices in all partition sets. It is not difficult to check that there are 14,400 such triangles. From this information, we compute $g(\Sigma) = 7,201$. Whence

$$7,201 \geq g(M_D) \geq 96.$$

The above approach applies to any Coxeter type situation. In case of the Davis manifold, improvements can be made by choosing smaller triangulations of the 120-cell that still have the desired partition properties. Our current best upper bound, however, remains at 5621, and does not improve the current gap in magnitudes. It would be interesting to see whether greater improvements can be obtained for the known hyperbolic 4-manifolds arising from Coxeter constructions.

Concluding remarks. The main challenge in obtaining lower bounds on trisection genus lies in dermining lower bounds on the genera of the 4-

dimensional 1-handlebodies. Mostow rigidity, of course, implies that every algebraic or topological invariant of a hyperbolic 4-manifold is a geometric invariant, but there is a philosophical distinction. It is in this spirit that we formulate the following:

Task 8. Determine stronger lower bounds on trisection genus of hyperbolic 4-manifolds using the *geometry* of the manifold.

Acknowledgements. The authors thank David Gay, Jonathan Hillman and Jeff Meier for interesting and helpful comments. This work is supported by ARC Future Fellowship FT170100316.

REFERENCES

- [1] AimPL, *Trisections and low-dimensional topology*. Available at <http://aimpl.org/trisections>.
- [2] R. Aranda, *Thin Position for 4-manifolds*. ArXiv:1805.08857.
- [3] M. Bell, J. Hass, J. Hyam Rubinstein and S. Tillmann, *Computing trisections of 4-manifolds*. Proc. Nat. Acad. Sci. USA **115** (2018), 43, 10901–10907.
- [4] N.A. Castro and B. Ozbagci, *Trisections of 4-manifolds via Lefschetz fibrations*. Math. Res. Lett. **26** (2019), 2, 383–420.
- [5] M.W. Davis, *A hyperbolic 4-manifold*. Proc. Amer. Math. Soc. **93** (1985), 2, 325–328.
- [6] D. Gay and R. Kirby, *Trisecting 4-manifolds*. Geom. Topol. **20** (2016), 6, 3097–3132.
- [7] M. Gromov, *Volume and bounded cohomology*. Publ. Math. Inst. Hautes Études Sci. **56** (1982), 5–99.
- [8] I.A. Grushko, *On the bases of a free product of groups*. Mat. Sb. **8** (1940), 169–182.
- [9] J. Hillman, *Four-Manifolds, Geometries and Knots*. Geometry and Topology Monographs **5**, Geometry and Topology Publications, Coventry, 2002.
- [10] D. Kotschick, *On the Gromov-Hitchin-Thorpe inequality*. C. R. Acad. Sci. Paris **326** (1998), 727–731.
- [11] M. Lackenby, *Heegaard splittings, the virtually Haken conjecture and property (τ)* . Invent. Math. **164** (2006), 2, 317–359.
- [12] W. Lück, *L^2 -torsion and 3-manifolds*. In: K. Johannson (Ed.), *Low Dimensional Topology*. Conf. Proc. Lect. Notes Geom. Topol. **3** (1994), International Press, Cambridge, 75–107.
- [13] J. Meier, *Trisections and spun 4-manifolds*. Math. Res. Lett. **25** (2018), 5, 1497–1524.
- [14] J. Meier, T. Schirmer and A. Zupan, *Classification of trisections and the generalized property R conjecture*. Proc. Amer. Math. Soc. **144** (2016), 11, 4983–4997.
- [15] J. Meier and A. Zupan, *Genus-two trisections are standard*. Geom. Topol. **21** (2017), 3, 1583–1630.
- [16] J. Milnor and W. Thurston, *Characteristic numbers of 3-manifolds*. Enseign. Math. (2) **23** (1977), 3–4, 249–254.
- [17] B.H. Neumann, *On the number of generators of a free product*. J. Lond. Math. Soc. **18** (1943), 12–20.

- [18] J.G. Ratcliffe and S.T. Tschantz, *On the Davis hyperbolic 4-manifold*. *Topology Appl.* **111** (2001), 3, 327–342.
- [19] J.H. Rubinstein and S. Tillmann, *Multisections of piecewise linear manifolds*. To appear in *Indiana Univ. Math. J.* <http://arxiv.org/abs/1602.03279>.
- [20] J. Spreer and S. Tillmann, *The trisection genus of the standard simply connected PL 4-manifolds*. 34th International Symposium on Computational Geometry (SoCG 2018). In: *Leibniz International Proceedings in Informatics (LIPICS)* **99** (2018), 71:1-71:13.

Received 11 March 2019

*University of California,
Department of Mathematics,
Santa Barbara, CA 93106, USA
mchu@math.ucsb.edu*

*University of Sydney,
School of Mathematics and Statistics F07,
NSW 2006 Australia
stephan.tillmann@sydney.edu.au*