

INSTITUTUL  
DE  
MATEMATICA

INSTITUTUL NATIONAL  
PENTRU CREATIE  
STIINTIFICA SI TEHNICA

---

ISSN 0250 3638

---

INDECOMPOSABLE COHEN MACAULAY MODULES  
AND THEIR MULTIPLICITIES

by

Dorin POPESCU

PREPRINT SERIES IN MATHEMATICS

No. 12/1988

---

BUCURESTI

Med 24810

INDECOMPOSABLE COHEN MACAULAY MODULES  
AND THEIR MULTIPLICITIES

by

Dorin POPESCU\*

February 1988

\*) Department of Mathematics, The National Institute for  
Scientific and Technical Creation, Bd. Păcii 220, 79622  
Bucharest, Romania.

# INDECOMPOSABLE COHEN-MACAULAY MODULES AND THEIR MULTIPLICITIES

Dorin Popescu

**ABSTRACT.** The main aim of this paper is to find a large class of rings for which there are indecomposable maximally Cohen-Macaulay modules of arbitrary high multiplicity (or rank in the case of domains).

## 1. Introduction

Let  $(A, \mathfrak{m})$  be a (commutative) henselian Cohen-Macaulay local ring and  $\text{CM}(A)$  the category of maximally Cohen-Macaulay  $A$ -modules (shortly MCM  $A$ -modules) i.e. of finitely generated modules  $M$  with  $\text{depth } M = \dim A$ . For  $s \in \mathbb{N}$  let  $n_A(s)$  be the cardinal of isomorphism classes of indecomposable modules  $M$  from  $\text{CM}(A)$  whose multiplicity  $e_A(M) = e(\mathfrak{m}, M) = s$ . Take  $n_A = \sum_{s \in \mathbb{N}} n_A(s)$ .

(1.1) First Brauer-Thrall type conjecture. If  $n_A = \infty$ , then  $n_A(s) \neq 0$  for infinitely many  $s$ .

When  $\dim A = 0$  then  $e_A(M) = \text{length}_A(M)$  and (1.1) holds by A. V. Roiter's theorem ([R], [Au<sub>1</sub>] or [P] (7.7)). Using the Auslander-Reiten theory for MCM modules (see [Au<sub>2</sub>], [P], [AR<sub>1</sub>], [Ya] or [P] Appendix) Y. Yoshino succeeded to solve positively (1.1) for reduced analytic algebras  $A$  over a perfect valued field  $k$  which are isolated singularities. Our Theorem (5.4) gives in particular the following

(1.2) Theorem. Let  $(A, \mathfrak{m})$  be a reduced excellent henselian local CM-ring,  $k := A/\mathfrak{m}$ ;  $p := \text{char } k$ . Suppose that

(i)  $\text{rk}_k(k^p) < \infty$  if  $p > 0$ ,

(ii)  $A$  is an isolated singularity,

(iii) if  $pA = 0$  then for every  $\mathfrak{q} \in \text{Spec } A$ ,  $\mathfrak{q} \neq \mathfrak{m}$  containing  $pA$ ,  $A_{\mathfrak{q}}/pA_{\mathfrak{q}}$  is regular.

Then (1.1) holds.

Note that (iii) follows from (ii) when  $A$  contains a field (i.e. the equal characteristic

---

1980 Mathematics Subject Classification. Primary 13H10, Secondary 13H15, 13J15; Key words and phrases. Maximally Cohen-Macaulay modules, Indecomposable modules, Multiplicity.



case). When  $A$  is a domain  $e_A(M) = e(A) \cdot \text{rank}(M)$  by  $[M_2]$  (14.8) so in the hypothesis of our Theorem there are indecomposable MCM-modules of arbitrary high rank if  $n_A = \infty$ . The proof follows entirely [Y] our contribution being mainly to extend his Lemmas (2.10), (2.12) in the following form (see (4.8)):

(1.3) Theorem. Let  $(A, \underline{m})$  be a reduced excellent henselian local CM - ring,  $k := A/\underline{m}$ ,  $p := \text{char } k$  and  $I_s(A)$  the ideal defining the singular locus of  $A$ , i.e.  $I_s(A) = \bigcap_{q \notin \text{Reg } A} q$ . Suppose that

i)  $[k : k^p] < \infty$  if  $p > 0$ ,

ii) if  $pA \neq 0$  then for every  $q \in \text{Reg } A$  containing  $pA$   $A_q/pA_q$  is regular,

iii)  $I_s(A) \subseteq \underline{m}$ , i.e.  $A$  is not regular.

Then there exists a positive integer  $r$  such that

1) A MCM  $A$ -module  $M$  is indecomposable iff  $M/I_s(A)^r M$  is indecomposable,

2) Two indecomposable MCM  $A$ -modules  $M, N$  are isomorphic iff  $M/I_s(A)^r M$  and  $N/I_s(A)^r N$  are isomorphic.

In particular this Theorem gives large classes of isolated singularities for which there exist Dieterich [D] reduction ideals.

In the hypothesis of (1.3) we get  $n_A \leq n_{\hat{A}}$  (see (4.10)) where  $\hat{A}$  is the completion of  $A$ . In particular we can improve the result from [K] and [BGS] for excellent henselian local rings (see (4.11)). Though (4.11) can be also obtained using the property of Artin approximation of excellent henselian local rings (see [Po] (1.3)) as we indicate in (5.6), we choose here an easier method (see §§ 3-4) which is entirely self contained and proves to be more powerful for these questions. Our Section 2 contains just preliminaries arranged more or less after [Y] which we include it here for the completeness. We supply here a proof of (2.5) because [S] was not available to us.

We would like to thank A. Brezuleanu and N. Radu for many helpful conversations on Theorem (4.4).

## 2. The singular locus of an excellent local ring.

Let  $A$  be an excellent ring. Then  $\text{Reg } A = \{q \in \text{Spec } A \mid A_q \text{ is regular}\}$  is an open set and  $I_s(A) = \bigcap_{q \notin \text{Reg } A} q$  defines the singular locus of  $A$ , i.e.

$$V(I_s(A)) = \text{Spec } A \setminus \text{Reg } A.$$

(2.1) Lemma. Let  $u: A \rightarrow B$  be a flat morphism of excellent rings. Then  $I_s(B) \subseteq \sqrt{u(I_s(A))}B$ .

Proof. If  $q \in \text{Reg } B$  then  $q \cap A \in \text{Reg } A$  by  $[M_1]$  (21.D). Thus a prime ideal



from  $B$  containing  $u(I_S(A))$  must contain also  $I_S(B)$ .  $\square$

(2.2) It will be useful also to express  $I_S(A)$  as the radical of a certain ideal of  $A$  whose elements can be precisely described. This is already well known for rings  $A$  which are essentially of finite type over a perfect field  $k$  because in that case the Jacobian criterion for smoothness [ $M_1$ ] (29.C) applies and we have  $I_S(A) = H_{A/k}$ . In general, given a finite presentation  $A$ -algebra  $B = A[X]/\underline{a}$ ,  $X = (X_1, \dots, X_n)$ , the nonsmooth locus of  $B$  over  $A$  is defined by the following ideal

$$H_{B/A} = \sqrt{\sum_f \Delta_f((f) : \underline{a})B}$$

where the sum is taken over all systems  $f$  of  $r$ -polynomials from  $\underline{a}$ ,  $r = 1, \dots, n$  being variable (see [Po] (2.1)). Using [Y] § 2 we will present such a description of  $I_S(A)$  when  $A$  is a Noetherian complete local ring having some additional properties.

(2.3) Till the end of this Section  $(R, m)$  is a reduced Noetherian complete local ring with a perfect residue field  $k$ . Then either  $R$  contains  $k$  or  $R$  is an algebra over a Cohen ring of residue field  $k$ , i.e. a complete DVR  $(T, t)$  which is an unramified extension of  $\mathbb{Z}_{(p)}$ ,  $p := \text{char } k > 0$ ,  $t := p \cdot 1 \in T$ . When  $R$  contains  $k$  we put  $T := k$  and  $t = 0$  in order to unify both situations.

Let  $\mathcal{R}(T, R)$  be the set of all prime ideals  $q \subseteq R$  for which  $T \rightarrow R_q$  is a regular morphism. Clearly  $\mathcal{R}(T, R) \subseteq \text{Reg } R$  because  $T$  is regular and regular morphisms preserve this property ([ $M_1$ ] (33.b)). When  $R$  contains  $k$  the other inclusion also holds,  $k$  being perfect. When  $R$  is in the unequal characteristic case ( $pR \neq 0$ ) then we suppose that

(\*)  $R_q/pR_q$  is regular for every  $q \in \text{Reg } R$

Thus in both situations we have  $\mathcal{R}(T, R) = \text{Reg } R$ .

(2.4) Let  $x = (x_1, \dots, x_n)$  be a system of elements from  $R$  such that  $(t, x)$  forms a system of parameters in  $R$ . From now on we suppose that  $R$  is a Cohen Macaulay ring (shortly a CM - ring). Then the canonical map  $T[[X]] \rightarrow R$ ,  $X = (X_1, \dots, X_n) \rightarrow x$  is finite and flat (hence free) by Cohen Structure Theorems and [ $M_1$ ] (36.B).

(2.5) Lemma (Scheja - Storch [S]). There exists  $x$  as above such that  $\text{ht}(H_{R/T[[x]]}) \geq 1$ , i.e. for every minimal prime ideal  $q \subseteq R$  the fraction field extension  $\text{Fr}(T[[x]]) \hookrightarrow R_q$  is (finite) separable.

**Proof.** When  $T \neq k$  there is nothing to show because  $\text{char } T = 0$ . Suppose  $T = k$ . Let  $q_1, \dots, q_s$  be the minimal prime ideals of  $R$  and take an arbitrary system of parameters  $y = (y_1, \dots, y_n)$  of  $R$ . If the field extensions  $\alpha_i : k((y)) \rightarrow R_{q_i}$ ,  $1 \leq i \leq s$  are

all separable then  $\text{ht}(H_{R/k[[y]]}) \geq 1$  by the Jacobian criterion for smoothness  $[M_1]$  (29.C). Suppose that  $(\alpha_i)_{1 \leq i \leq e}$  are not separable for a certain  $e$ ,  $1 \leq e \leq s$ . Then  $p > 0$  and for every  $i$ ,  $1 \leq i \leq e$  there exists an element  $z_i \in R_{q_i} \setminus k((y))$  such that  $z_i^p \in k((y))$ . Since  $\alpha_i$  is finite we have

$$k((y)) \otimes_{k[[y]]} R \cong \prod_{i=1}^s R_{q_i}.$$

Thus we can find one  $z \in R$  and  $w \in k[[y]]$  such that  $z/w$  corresponds to  $(z_1, \dots, z_e, y_n, \dots, y_n)$  by the above isomorphism. Then  $h := z^p \in k[[y]]$  and  $z \in k[[y]]$ . Adding a constant to  $z$  we can suppose that  $h \in (y)k[[y]]$ . If  $h \in k[[y^p]]$  then  $h \in k^p[[y^p]]$  ( $k$  is perfect) and so  $z \in k[[y]]$  which is not possible.

Suppose that  $h \notin k[[y_1, \dots, y_{n-1}, y_n^p]]$ . After a coordinate transformation we can suppose also that  $h$  is regular in  $y_n$ . Applying Weierstrass Preparation Theorem for  $U - h$  in  $k[[y, U]]$  we find a distinguished polynomial

$$(1) P = y_n^r + \sum_{i=1}^r a_i y_n^{r-i}, \quad a_i \in k[[y_1, \dots, y_{n-1}, U]], \quad a_i(0) = 0$$

and an invertible formal power series  $g \in k[[y, U]]$  such that

$$(2) U - h = Pg$$

Substituting  $U = h$  in  $P$  we get

$$(3) y_n^r + \sum_{i=1}^r a_i(y_1, \dots, y_{n-1}, h) y_n^{r-i} = 0$$

because  $g(U = h) \neq 0$  since  $g(0) \neq 0$  and  $h(0) = 0$ . Applying  $\partial / \partial y_n$  in (2) we obtain

$$(\partial P / \partial y_n)g + P(\partial g / \partial y_n) = -\partial h / \partial y_n \neq 0$$

and substituting  $U = h$  we get  $(\partial P / \partial y_n)(U = h) \neq 0$ . Thus (3) defines a separable equation for  $y_n$  over  $k[[y_1, \dots, y_{n-1}, z^p]]$ . In particular  $y_n$  is separable over  $S := k[[y_1, \dots, y_{n-1}, z]]$ . Denote  $y' = (y_1, \dots, y_{n-1}, z)$ . We have

$$[R_{q_i} : k((y'))]_{\text{ins}} = [R_{q_i} : k((y))]_{\text{ins}} - p$$

for every  $i = 1, \dots, e$ , where  $[ ]_{\text{ins}}$  denotes the inseparable degree. Repeating this procedure inductively we finally find a system of parameters  $x$  in  $R$  such that  $k((x)) \hookrightarrow R_{q_i}$  is separable for every  $i$ .  $\square$



(2.6) Remark. If  $k$  is not perfect then the above Lemma doesn't hold. If  $a \in k \setminus k^p$  then  $A = k[[X, Y]]/(X^p + aY^p)$  (after  $[Y]$  (2.7)) is a counterexample.

(2.7) Lemma. Let  $q \in \text{Reg } R$ . Then there exists a system of elements  $x$  in  $R$  such that

(i)  $(t, x)$  is a system of parameters in  $R$ ,

(ii)  $H_{R/T[[x]]} \not\subset q$ .

Proof. If  $t = 0$  then we choose a system of elements  $y$  in  $R$  which forms in  $R_q$  a regular system of parameters. If  $t \in q$  then by condition (2.3) (\*) we get  $R_q/tR_q$  regular. Thus there exists  $y$  such that  $(t, y)$  form in  $R_q$  a regular system of parameters. By Lemma (2.5) there exists a system of elements  $z$  in  $R$  which forms a system of parameters  $\bar{z}$  in  $R/\underline{a}$ ,  $\underline{a} := \sqrt{(t, y)}$  such that the map  $(T/T \cap \underline{a})[[\bar{z}]] \rightarrow R/\underline{a}$  is generically smooth. (Note that  $R/(t, z)$  is CM (see  $[M_1]$  (16.C)) and so  $R/\underline{a}$  is CM too). Since  $q$  is a minimal prime ideal containing  $\underline{a}$  we get  $(T/T \cap q)[[\bar{z}]] \rightarrow R/q$  separable and so the map  $T[[y, z]] \rightarrow R_q$  is etale. Thus  $x = (y, z)$  works.

Suppose now  $t \notin q$  then as above we can choose  $y$  in  $R$  which forms a regular system of parameters in  $R_q$ . Take a system of elements  $z$  in  $R$  such that  $(t, z)$  forms modulo  $q$  a system of parameters in  $R/q$ . Then  $(t, y, z)$  forms a system of parameters in  $R$  and  $T[[y, z]] \rightarrow R_q$  is etale ( $\text{char } R/q = 0$ ). Thus  $x = (y, z)$  works.  $\square$

(2.8) Corollary.  $I_S(R) = \sqrt{\sum_x H_{R/T[[x]]}}$ , where the sum is taken over all systems of elements  $x$  such that  $(t, x)$  forms a system of parameters of  $R$ .

Proof. If  $q \in \text{Spec } R$  does not contain  $H_{R/T[[x]]}$  for a certain system  $x$  then the map  $T[[x]] \rightarrow R_q$  is etale and so  $R_q$  is regular because  $T[[x]]$  is so. Conversely if  $q \in \text{Reg } R$  then by Lemma (2.7) there exists  $x$  such that  $q \not\supset H_{R/T[[x]]}$ .  $\square$

(2.9) Let  $S \subseteq R$  be a regular local subring such that  $R$  is a finitely generated free  $S$ -module,  $R^e := R \otimes_S R$  the enveloping algebra of  $R$  over  $S$  and  $\mu: R^e \rightarrow R$  the multiplication map. Denote  $I = \text{Ker } \mu$ . The ideal  $\mathcal{N}_S^R = \mu(\text{Ann}_{R^e} I)$  is called the Noether different of  $R$  over  $S$ .

(2.10) Lemma.  $\mathcal{N}_S^R \cdot \Omega_{R/S} = 0$  and  $H_{R/S} = \sqrt{\mathcal{N}_S^R}$ .

Proof. The first equality is trivial because  $\Omega_{R/S} = I/I^2$ . Let  $q \in R$  be a prime ideal. If  $q \not\supset \mathcal{N}_S^R$  then  $\Omega_{R_q/S} = \Omega_{R/S} \otimes_R R_q = 0$  as above. Since  $S \subseteq R$  is finite free we get  $S \rightarrow R_q$  etale, i.e.  $q \not\supset H_{R/S}$ . Conversely if  $q \not\supset H_{R/S}$  then  $S \rightarrow R_q$  is etale and so



$\Omega_{R/S} \otimes_R R_Q = 0$ . Thus  $I_Q = I_Q^2$  for a certain prime ideal  $Q \subseteq R^e$ ,  $Q \supseteq I$  such that  $\mu(Q) = q$ .

By Nakayama Lemma we get  $I_Q = 0$  and so  $Q \not\subseteq \text{Ann}_{R^e} I$ . Thus  $\mathcal{N}_S^R \not\subseteq q$ .

(2.11) We end this Section by listing some facts from Hochschild cohomology, which can be found in [P] Ch.11. Let  $B \subseteq A$  be an extension of rings. The  $n$ 'th Hochschild cohomology functors  $H_B^n(A, -)$ ,  $n \geq 0$  are defined on the category of  $A$ -bimodules with values in the category of  $A$ -modules and have the following properties

i)  $H_B^0(A, M) = M^{(A)} := \{x \in M \mid ax = xa \text{ for every } a \in A\}$  for all  $A$ -bimodules  $M$ ,

ii) If  $M, N$  are two  $A$ -modules then  $\text{Hom}_B(M, N)$  is an  $A$ -bimodule [the left (resp. right) action of  $A$  on  $\text{Hom}_B(M, N)$  is given as the one induced from the action on  $N$  (resp.  $M$ )] and  $H_B^0(A, \text{Hom}_B(M, N)) = \text{Hom}_A(M, N)$ .

iii)  $H_B^1(A, M)$  is a factor  $A$ -module of  $\text{Der}_B(A, M) = \text{Hom}_A(\Omega_{A/B}, M)$

iv) If  $A$  is a projective module over  $B$  and

$$0 \longrightarrow M' \longrightarrow M \longrightarrow M'' \longrightarrow 0$$

is a short exact sequence of  $A$ -bimodules then there exist some  $A$ -morphisms  $\partial^{(n)} : H_B^n(A, M'') \longrightarrow H_B^{n+1}(A, M')$ ,  $n \geq 0$  such that the following sequence is exact.

$$\begin{aligned} 0 \longrightarrow H_B^0(A, M') \longrightarrow H_B^0(A, M) \longrightarrow H_B^0(A, M'') \longrightarrow H_B^1(A, M') \longrightarrow \dots \\ \dots \longrightarrow H_B^n(A, M') \longrightarrow H_B^n(A, M) \longrightarrow H_B^n(A, M'') \longrightarrow H_B^{n+1}(A, M') \longrightarrow \dots \end{aligned}$$

(2.12) Lemma. Let  $S \subseteq R$  be as in (2.9) and  $M$  an  $R$ -bimodule. Then  $\mathcal{N}_S^R \cdot H_S^1(R, M) = 0$ .

Proof. By Lemma (2.10) we have  $\mathcal{N}_S^R \Omega_{R/S} = 0$  and so  $\mathcal{N}_S^R \cdot \text{Hom}_R(\Omega_{R/S}, M) = 0$ .

Now apply (2.11) iii).  $\square$

### 3. CM - approximation

(3.1) Lemma. Let  $S \subseteq R$  be an extension of Noetherian rings such that  $R$  is a finitely generated projective module over  $S$ ,  $x$  an element from  $\mathcal{N}_S^R$  and  $M, N$  two finitely generated  $R$ -modules such that  $M$  is projective over  $S$ . Let  $e \in \mathbb{N}$  be a positive integer such that  $\text{Ann}_N x^e = \{z \in N \mid x^e z = 0\} = \text{Ann}_N x^{e+1}$  and  $s \in \mathbb{N}$ . Then for every linear  $R$ -map  $\varphi : M \longrightarrow N/x^{e+s+1}N$  there exists a linear  $R$ -map  $\psi : M \longrightarrow N$  which makes commutative the following diagram

$$\begin{array}{ccc}
 M & \xrightarrow{\varphi} & N/x^{e+s+1}N \\
 \downarrow \psi & & \downarrow \\
 N & \xrightarrow{\quad} & N/x^{e+s}N
 \end{array}$$

**Proof.** Let  $N' := \text{Ann}_N x^e$ . We have the following commutative diagram:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & N/N' & \xrightarrow{x^{e+s+1}} & N/N' & \longrightarrow & N/N' + x^{e+s+1}N \longrightarrow 0 \\
 & & \downarrow x & & \downarrow & & \downarrow \\
 0 & \longrightarrow & N/N' & \xrightarrow{x^{e+s}} & N/N' & \longrightarrow & N/N' + x^{e+s}N \longrightarrow 0
 \end{array}
 \quad (1)$$

in which the bases are exact. Indeed if  $x^{e+s}z \in N'$  for a certain  $z \in N$  then  $x^{2e+s}z = 0$  and so  $z \in \text{Ann}_N x^{2e+s} = N'$ . Applying the functor  $\text{Hom}_S(M, -)$  to (1) we get the following commutative diagram:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \text{Hom}_S(M, N/N') & \longrightarrow & \text{Hom}_S(M, N/N') & \longrightarrow & \text{Hom}_S(M, N/N' + x^{e+s+1}N) \longrightarrow 0 \\
 & & \downarrow x & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \text{Hom}_S(M, N/N') & \longrightarrow & \text{Hom}_S(M, N/N') & \longrightarrow & \text{Hom}_S(M, N/N' + x^{e+s}N) \longrightarrow 0
 \end{array}
 \quad (2)$$

where the bases are exact because  $M$  is projective over  $S$ . Clearly these bases are also exact sequences of  $R$ -bimodules and applying the Hochschild cohomology functors we get the following commutative diagram (see (2.11) ii)):

$$\begin{array}{ccccccc}
 \text{Hom}_R(M, N/N') & \longrightarrow & \text{Hom}_R(M, N/N' + x^{e+s+1}N) & \longrightarrow & H_S^1(R, \text{Hom}_S(M, N/N')) \\
 \downarrow & & \downarrow & & \downarrow x \\
 \text{Hom}_R(M, N/N') & \longrightarrow & \text{Hom}_R(M, N/N' + x^{e+s}N) & \longrightarrow & H_S^1(R, \text{Hom}_S(M, N/N'))
 \end{array}
 \quad (3)$$

in which the bases are exact (see (2.11) iv)). Since the last vertical map is zero by Lemma (2.12) we get a linear  $R$ -map  $\alpha : M \longrightarrow N/N'$  such that the following diagram is commutative



$$(4) \quad \begin{array}{ccccc} M & \xrightarrow{\psi} & N/x^{e+s+1}N & \longrightarrow & N/N' + x^{e+s+1}N \\ \downarrow \alpha & & & & \downarrow \\ N/N' & \longrightarrow & & & N/N' + x^{e+s}N \end{array}$$

Note that in the following diagram

$$(5) \quad \begin{array}{ccccc} M & \xrightarrow{\quad} & N/x^{e+s+1}N & & \\ \parallel & & \downarrow & & \\ M & \xrightarrow{\psi} & N/N' \cap x^{e+s}N & \longrightarrow & N/x^{e+s}N \\ \parallel & & \downarrow & & \downarrow \\ M & \xrightarrow{\alpha} & N/N' & \longrightarrow & N/N' + x^{e+s}N \end{array}$$

the small square is cartesian and so there exists  $\psi$  which makes (5) commutative.

Remains to show that  $N' \cap x^{e+s}N = 0$ . Indeed let  $y \in N' \cap x^{e+s}N$  and  $z \in N$  with  $y = x^{e+s}z$ . Then  $0 = x^e y = x^{2e+s}z$  and so  $z \in N'$ , i.e.  $y = x^{e+s}z = 0$ .  $\square$

Remark. Roughly speaking Lemma (3.1) says that given  $M, N$  there exists a function  $\gamma : \mathbb{N} \rightarrow \mathbb{N}$  such that every linear  $R$ -map  $\varphi : M \rightarrow N/x^{\gamma(s)}N$ ,  $s \in \mathbb{N}$  can be lifted to a linear  $R$ -map  $\psi : M \rightarrow N$  such that  $(R/x^s R) \otimes_R \psi = (R/x^s R) \otimes_R \varphi$ . But this follows easily from a linear form of the strong approximation theorem (see [Po] § 1) which holds in fact in every Noetherian local ring  $R$  for every element  $x \in R$ . Thus the importance of Lemma (3.1) consists just in giving to  $\gamma$  a precised form.

(3.2) Lemma. Let  $B \subset A$  be a finite flat extension of Noetherian rings,  $\underline{a} \subset A$  an ideal and  $x \in H_{A/B}$  an element. Then there exists a positive integer  $r$  such that for every finitely generated  $A$ -module  $N$  which is free over  $B$  it holds

$$(\underline{a}N : x^r)_N = (\underline{a}N : x^{r+1})_N,$$

$$\text{where } (\underline{a}N : x^r)_N = \{ z \in N \mid x^r z \in \underline{a}N \}.$$

Proof. Step 1 Reduction to the case  $(\underline{a} : x) = \underline{a}$ .

Since  $A$  is Noetherian we have  $\underline{a}' := (\underline{a} : x^n) = (\underline{a} : x^{n+1})$  for a certain positive integer  $n$ . If  $xy \in \underline{a}'$  for a certain  $y \in A$  then  $x^{n+1}y \in \underline{a}$  and so  $y \in \underline{a}'$ , i.e.  $(\underline{a}' : x) = \underline{a}'$ .



Suppose that  $r' \in \mathbb{N}$  satisfies our Lemma for  $x$  and  $\underline{a}'$ . Then  $r = n + r'$  works. Indeed, let  $N$  be as in our Lemma. If  $x^s z \in \underline{a}N \subseteq \underline{a}'N$  for some  $s \in \mathbb{N}$  and  $z \in N$  then  $x^{r'} z \in \underline{a}'N$  because  $(\underline{a}'N : x^{r'})_N = (\underline{a}'N : x^{r'+1})_N$ . Thus  $x^{r'} z \in x^n \underline{a}'N \subseteq \underline{a}N$ .

Remark.  $\text{Ass}_A(A/\underline{a}') = \{q \in \text{Ass}_A(A/\underline{a}) \mid x \notin q\}$ .

Let  $\underline{a} = \bigcap_{i=1}^e Q_i$  be an irredundant prime decomposition of  $\underline{a}$ ,  $q_i := \sqrt{Q_i}$ ,  $q'_i := q_i \cap B$ ,  $Q'_i := Q_i \cap B$ ,  $\underline{b} := \underline{a} \cap B = \bigcap_{i=1}^e Q'_i$  and  $k'_i \subseteq k_i$  the residue field extension of  $B_{q'_i} \subset A_{q_i}$ .

Step 2. Case when  $k'_i = k_i$ ,  $1 \leq i \leq e$ .

By Step 1 we may suppose that  $(\underline{a}:x) = \underline{a}$ . Fix an  $i$ ,  $1 \leq i \leq e$ . Clearly  $x \notin q_i$  because  $x$  is a nonzero divisor of  $A/\underline{a}$ . Then the map  $B_{q'_i} \rightarrow A_{q_i}$  is etale and so  $q_i A_{q_i} = q'_i A_{q_i}$ . Since  $k'_i = k_i$  the extension  $B_{q'_i} \subset A_{q_i}$  is dense. In particular we have

$$B_{q'_i}/Q'_i B_{q'_i} \cong A_{q_i}/Q'_i A_{q_i}$$

and it follows  $Q'_i A_{q_i} = Q_i A_{q_i}$ .

We show that  $r = 0$  satisfies this case. Let  $N$  be as in our Lemma, and  $z \in N$  such that  $xz \in \underline{a}N$ . Then  $z \in Q_1 N_{q_1} = Q'_1 N_{q_1}$ . Thus there exists an element  $y_1 \in A \setminus q_1$  such that  $y_1 z \in Q'_1 N$ . Since  $B/q'_1 \rightarrow A/q_1$  is finite we get  $(y_1 A) \cap (B \setminus q'_1) \neq \emptyset$ . Thus changing  $y_1$  by one of its multiple we may suppose that  $y_1 \in B \setminus q'_1$ , i.e.  $z \in Q'_1 N_{q'_1}$ . Since  $N$  is free over  $B$  we have

$$\underline{b}N = \bigcap_{j=1}^e Q'_j N$$

and  $Q'_j N$  is exactly the  $q'_j$ -primary submodule of  $N$  associated to  $\underline{b}N$ . Then

$$N \cap Q'_j N_{q'_j} = Q'_j N$$

and so

$$z \in N \cap \left( \bigcap_{j=1}^e Q'_j N_{q'_j} \right) = \underline{b}N \subseteq \underline{a}N.$$

Step 3. Case when there exists a faithfully flat  $B$ -algebra  $C$  such that for every prime ideal  $q$  associated to  $\underline{C} \otimes_B \underline{a}$  in  $D := C \otimes_B A$  the residue field extension of  $C_q \cap C \hookrightarrow D_q$  is trivial.

We apply Step 2 to the case  $C \subseteq D$ ,  $\underline{a}D$ ,  $x' = 1 \otimes x \in D$ . Clearly

$x' \in D \otimes_A H_{A/B} \subseteq H_{D/C}$ . Then there exists  $r$  such that for every finitely generated  $D$ -module  $N'$  which is free over  $C$  it follows

$$(\underline{a}N' : x'^r)_{N'} = (\underline{a}N' : x'^{r+1})_{N'}$$

Let  $N$  be a finitely generated  $A$ -module which is free over  $B$  and take  $N'' = D \otimes_A N$ . Then  $N''$  is free over  $C$  and so we get in particular

$$(\underline{a}N'' : x'^r)_{N''} = (\underline{a}N'' : x'^{r+1})_{N''}$$

But  $(\underline{a}N'' : x'^r)_{N''} = D \otimes_A (\underline{a}N : x'^r)_N$ . Indeed,  $(\underline{a}N : x'^r)_N$  is exactly the kernel of the composed map  $f : N \xrightarrow{x'^r} N \longrightarrow N/\underline{a}N$  and by flatness  $\text{Ker}(D \otimes_A f) = D \otimes_A \text{Ker} f$ . Thus the inclusion  $u : (\underline{a}N : x'^r)_N \hookrightarrow (\underline{a}N : x'^{r+1})_N$  goes by base change in an equality. Since  $D$  is a faithfully flat  $A$ -algebra we get  $u$  surjective too.

#### Step 4 General case - reduction to Step 3

We need the following

(3.3) Lemma. Let  $S \subseteq R$  be a finite flat extension of Noetherian rings and denote

$$d_{R/S}'' = \max_{q' \in \text{Spec } S} \sum_{\substack{q \in \text{Spec } R \\ q \cap S = q'}} ([k(q) : k(q')] - 1),$$

where  $k(q)$  denotes the residue field of  $R_q$ . Then  $d_{R/S} < \infty$  and  $d_{R \otimes_S R/R} < d_{R/S}$  if  $d_{R/S} > 0$ , where the structural map  $R \longrightarrow R \otimes_S R$  is given by  $y \longrightarrow y \otimes 1$ .

Applying by recurrence the above Lemma we get finally a finite flat  $B$ -algebra  $C$  of the form  $A \otimes_B A \otimes \dots \otimes_B A$  such that  $d_{C \otimes_B A/C} = 0$  i.e.  $k(q \cap C) = k(q)$  for all  $q \in \text{Spec}(C \otimes_B A)$ . Since a finite flat extension is faithfully flat we are ready.  $\square$

Proof of Lemma (3.3). Let  $q' \in \text{Spec } S$ . Then

$$d_{R/S, q'} = \sum_{\substack{q \in \text{Spec } R \\ q \cap S = q'}} ([k(q) : k(q')] - 1) < \text{rank}_{k(q')} k(q') \otimes_S R,$$

the last number being bounded by the minimal number of generators of  $R$  over  $S$ . It is enough to show that

$$d_{R \otimes_S R/R, q} < d_{R/S, q'}$$



for every  $q \in \text{Spec } R$  lying over  $q'$  and such that  $d_{R/S, q'} > 0$ . So by base change we reduce the question to the case when  $S = k(q') =: k$ . Then  $R$  is Artinian. Let  $(k_i)_{1 \leq i \leq e}$  be its residue fields. It is enough to show that

$$d_{k_1 \otimes_k k_i / k} \leq d_{k_i / k}, \quad 1 \leq i \leq e \text{ and } d_{k_1 \otimes_k k_i} < d_{k_i / k} \text{ if } k \neq k_1.$$

First inequality is clear because

$$1 + d_{k_1 \otimes_k k_i / k} \leq \text{rank}_{k_1} k_1 \otimes_k k_i = \text{rank}_k k_i = d_{k_i / k} + 1$$

The equality holds only when  $k_1 \otimes_k k_i$  is a field. But  $k_1 \otimes_k k_i$  is not a field so the second inequality holds too.  $\square$

(3.4) Lemma. Let  $S \subseteq R$  be an extension of Noetherian rings such that  $R$  is a finitely generated projective module over  $S$ ,  $x$  an element from  $\mathcal{A}_S^R$  and  $\underline{a} \subseteq R$  an ideal. Then there exists an increasing function  $\gamma: \mathbb{N} \rightarrow \mathbb{N}$  such that for every  $s \in \mathbb{N}$ , for every finitely generated  $R$ -modules  $M, N$  which are free over  $S$  and for every linear  $R$ -map  $\varphi: M \rightarrow N/(\underline{a}, x^{\gamma(s)})N$  there exists a linear  $R$ -map  $\psi: M \rightarrow N/\underline{a}N$  which makes commutative the following diagram:

$$\begin{array}{ccc} M & \xrightarrow{\varphi} & N/(\underline{a}, x^{\gamma(s)})N \\ \downarrow & & \downarrow \\ N/\underline{a}N & \xrightarrow{\psi} & N/(\underline{a}, x^s)N \end{array}$$

Proof. Let  $r$  be the integer given by Lemma (3.3) for  $x$  and  $\underline{a}$ . Define  $\gamma$  by  $\gamma(s) = 1 + \max\{r, s\}$ . Then given  $M, N, s, \varphi$  as in our Lemma we find the wanted  $\psi$  applying Lemma (3.1) for  $x$ ,  $\bar{N} = N/\underline{a}N$  and  $e = r$ .  $\square$

(3.4) Lemma. Let  $x = (x_1, \dots, x_n)$  be a system of elements from a Noetherian ring  $R$  such that for every  $i$ ,  $1 \leq i \leq n$  there exists a Noetherian subring  $S_i$  of  $R$  such that

i)  $R$  is finite free over  $S_i$ ,

ii)  $x_i \in \mathcal{A}_{S_i}^R$ .

Then there exists an increasing function  $\gamma: \mathbb{N} \rightarrow \mathbb{N}$  such that for every  $s \in \mathbb{N}$ , every finitely generated  $R$ -modules  $M, N$  which are free over all  $(S_i)_{1 \leq i \leq n}$  and for every linear  $R$ -map  $\varphi: M \rightarrow N/x^{\gamma(s)}N$  there exists a linear  $R$ -map  $\psi: M \rightarrow N$  which makes



commutative the following diagram:

$$\begin{array}{ccc}
 M & \xrightarrow{\varphi} & N/x^{\vartheta(s)}N \\
 \downarrow \Psi & & \downarrow \\
 N & \xrightarrow{\quad} & N/x^sN
 \end{array}
 \quad (*)$$

**Proof.** Denote  $\underline{b}_i = (x_1, \dots, x_i)$ ,  $i = 1, \dots, n$ . Apply induction on  $n$ . If  $n = 1$  then apply Lemma (3.4) for  $x_1$  and  $\underline{a} = 0$ . Suppose now that it is given a function  $\vartheta'$  which works for  $\underline{b}_{n-1}$ . Let  $s \in N$  and  $\vartheta''_s$  be the function given by Lemma (3.4) for  $x_n$  and  $\underline{a} = \underline{b}_{n-1}$ . Define  $\vartheta: N \rightarrow N$  by  $\vartheta(s) = \vartheta'(s) + \vartheta''_s(s)$ . Let  $M, N$  be two finitely generated  $R$ -modules which are free over all  $(S_i)_{1 \leq i \leq n}$  and  $\varphi: M \rightarrow N/\underline{b}_n^{\vartheta(s)}N$  a linear  $R$ -map. Then there exists a linear  $R$ -map  $\alpha: \bar{M} \rightarrow \bar{N} := N/\underline{b}_{n-1}^{\vartheta'(s)}N$  which makes commutative the following diagram:

$$\begin{array}{ccc}
 M & \xrightarrow{\varphi} & N/\underline{b}_n^{\vartheta(s)}N \\
 \downarrow \alpha & & \downarrow \\
 \bar{M} & \xrightarrow{\quad} & \bar{N}/x_n^{\vartheta''_s(s)}\bar{N} \cong N/(\underline{b}_{n-1}^{\vartheta'(s)}, x_n^{\vartheta''_s(s)})N \\
 \downarrow & & \downarrow \\
 \bar{N} & \xrightarrow{\quad} & \bar{N}/x_n^s\bar{N} \cong N/(\underline{b}_{n-1}^{\vartheta'(s)}, x_n^s)N
 \end{array}$$

Thus there exists a linear  $R$ -map  $\psi: M \rightarrow N$  which makes commutative the following diagram

$$\begin{array}{ccccc}
 M & \xrightarrow{\alpha} & \bar{N} \cong N/\underline{b}_{n-1}^{\vartheta'(s)}N & \xrightarrow{\quad} & N/(\underline{b}_{n-1}^{\vartheta'(s)}, x_n^s)N \\
 \downarrow \psi & & \downarrow & & \downarrow \\
 N & \xrightarrow{\quad} & N/\underline{b}_{n-1}^sN & \xrightarrow{\quad} & N/\underline{b}_n^sN
 \end{array}$$

Clearly  $\psi$  makes also  $(*)$  commutative.  $\square$

(3.6) Let  $A$  be a CM local ring and  $M$  a MCM  $A$ -module. Then every system of parameters from  $A$  is a  $M$ -regular sequence. Let  $\underline{a} \subset A$  be a proper ideal.

The couple  $(A, \underline{a})$  is a CM - approximation if there exists a function  $\vartheta: N \rightarrow N$  (called CM - function) such that for every  $s \in N$ , every two MCM  $R$ -modules  $M, N$  and every linear  $R$ -map  $\varphi: M \rightarrow N/\underline{a}^{\vartheta(s)}N$  there exists a linear  $R$ -map  $\psi: M \rightarrow N$  such that  $(A/\underline{a}^s)_A \varphi \cong (A/\underline{a}^s)_A \psi$  in other words the following diagram is commutative:

$$\begin{array}{ccc}
 M & \xrightarrow{\quad} & N/a^{\vartheta(s)}N \\
 \vdots & & \downarrow \\
 N & \xrightarrow{\quad} & N/a^sN
 \end{array}$$

(3.7) Proposition. Let  $(R, \underline{m})$  be a reduced complete local CM-ring with a perfect residue field  $k$ ,  $p := \text{char } k$  and  $I_s(R)$  the ideal defining the singular locus of  $R$ . Suppose that for every  $q \in \text{Reg } R$  containing  $pR$  the ring  $R_q/pR_q$  is regular and  $I_s(R) \subseteq \underline{m}$ . Then  $(R, I_s(R))$  is a CM-approximation.

Proof. Let  $T \subseteq R$  be the Cohen ring of residue field  $k$  (see (2.3)). By Lemma (2.10) and Corollary (2.8) we have

$$I_s(R) = \sqrt{\sum_x \mathcal{N}_T^R[[x]]},$$

where the sum is taken over all systems of elements  $x$  such that  $(t, x)$  forms a system of parameters of  $R$ . Then we can find a system of elements  $y = (y_1, \dots, y_r)$  in  $I_s(R)$  such that

- 1)  $I_s(R) = \sqrt{yR}$
- 2) for every  $i = 1, \dots, r$  there exists a system of elements  $x^{(i)}$  of  $R$  such that  $(t, x^{(i)})$  forms a system of parameters of  $R$  and  $y_i \in \mathcal{N}_T^R[[x^{(i)}]]$ .

Since  $R$  is CM the inclusion  $S_i := T[[x^{(i)}]] \subset R$  is finite flat (so free). Let  $\vartheta': N \rightarrow N$  be the function given by Lemma (3.5) for  $y$ . If  $M, N$  are two MCM  $R$ -modules then  $(t, x^{(i)})$  is a regular  $M$  or  $N$ -sequence for all  $i$ . Thus  $M$  and  $N$  are finitely generated flat over  $S_i$ ,  $1 \leq i \leq r$  (see  $[M_1]$  (20.C)) and so free.

Now let  $u$  be a positive integer such that  $I_s(R)^u \subset yR$  and note that  $\vartheta$  given by  $\vartheta(s) = u \vartheta'(s)$  works.  $\square$

#### 4. CM - reduction ideals

(4.1) Lemma. Let  $(A, \underline{m})$  be a Noetherian local ring and  $\underline{a} \subset A$  an ideal. The following statements are equivalent:

- i)  $(A, \underline{a})$  is a CM-approximation,
- ii)  $(A, \sqrt{\underline{a}})$  is a CM-approximation.

Proof. Let  $u$  be a positive integer such that  $(\sqrt{\underline{a}})^u \subset \underline{a}$ . If i) holds and  $\vartheta: N \rightarrow N$  is the associated CM-function then as in the proof of Proposition (3.7) the function  $\overline{\vartheta}$  given by  $\overline{\vartheta}(s) = u \vartheta(s)$  works for  $(A, \sqrt{\underline{a}})$ . If ii) holds and  $\overline{\vartheta}$  is the associated CM-function then the function  $\vartheta$  given by  $\vartheta(s) = \overline{\vartheta}(su)$  works. Indeed, let  $M, N$  be two



MCM  $A$ -modules,  $s \in \mathbb{N}$  and  $\varphi: M \rightarrow N/\underline{a}^{\gamma(s)}N$  a linear  $A$ -map then there exists a linear map  $\psi: M \rightarrow N$  such that the following diagram commutes:

$$\begin{array}{ccccccc}
 M & \xrightarrow{\varphi} & N/\underline{a}^{\gamma(su)}N & \longrightarrow & N/(\sqrt{a})^{\gamma(su)}N & \longrightarrow & N/\underline{a}^sN \\
 \psi \downarrow & & & & \downarrow & & \parallel \\
 N & \longrightarrow & N/(\sqrt{a})^{su}N & \longrightarrow & N/\underline{a}^sN & & 
 \end{array}$$

(4.2) Lemma. Let  $A \rightarrow B$  be a flat local morphism of CM-local rings and  $\underline{a} \subset A$  an ideal. If  $(B, \underline{a}B)$  is a CM-approximation then  $(A, \underline{a})$  is too.

Proof. We claim that the CM-function  $\gamma$  associated to  $(B, \underline{a}B)$  works also for  $(A, \underline{a})$ . Indeed, let  $M, N$  be two MCM  $A$ -modules,  $s \in \mathbb{N}$  and  $\varphi: M \rightarrow N/\underline{a}^{\gamma(s)}N$  a linear  $A$ -map. Then  $\bar{M} = B \otimes_A M$ ,  $\bar{N} = B \otimes_A N$  are MCM  $B$ -modules since by flatness  $\text{depth } \bar{M} = \text{depth}_A M + \text{depth}(B/\underline{m}B) = \text{depth } A + \text{depth}(B/\underline{m}B) = \text{depth } B$  where  $\underline{m}$  denotes the maximal ideal of  $A$ . Thus there exists a linear  $B$ -map  $\bar{\psi}: \bar{M} \rightarrow \bar{N}$  such that the following diagram commutes

$$\begin{array}{ccc}
 \bar{M} & \xrightarrow{B \otimes_A \varphi} & \bar{N}/\underline{a}^{\gamma(s)}\bar{N} \\
 \bar{\psi} \downarrow & & \downarrow \\
 \bar{N} & \longrightarrow & \bar{N}/\underline{a}^s\bar{N}
 \end{array}$$

Since  $M, N$  are finitely generated modules, the existence of  $\bar{\psi}: \bar{M} \rightarrow \bar{N}$  such that the above diagram commutes means in other words that a certain linear system of equations  $L$  over  $A$  has a solution in  $B$ . Indeed, let  $M = A^n/(z_1, \dots, z_e)$ ,  $z_i = (z_{ij})_{1 \leq j \leq n}$ ,  $N = A^{n'}/(z'_1, \dots, z'_{e'})$ ,  $z'_\lambda = (z'_{\lambda\mu})_{1 \leq \mu \leq n'}$ ,  $\underline{a} = (a_1, \dots, a_v)$  a system of generators of  $\underline{a}$  and  $\varphi$  is given by the matrix  $(w_{j\mu})_{\substack{1 \leq j \leq n \\ 1 \leq \mu \leq n'}}$ . Then  $L$  has the following form

$$\sum_{j=1}^n z_{ij} X_{j\mu} = \sum_{\lambda=1}^{e'} Y_{i\lambda} z'_{\lambda\mu}, \quad 1 \leq i \leq e, \quad 1 \leq \mu \leq n'$$

$$X_{j\mu} - w_{j\mu} = \sum_{\alpha=1}^v a_\alpha U_{\alpha j\mu} + \sum_{\lambda=1}^{e'} Y'_{j\lambda} z'_{\lambda\mu}, \quad 1 \leq j \leq n$$

Clearly  $\bar{\psi}$  gives a solution of  $L$  in  $B$ . By faithfully flatness  $L$  has also a solution  $(x_{j\mu}, y_{i\lambda}, y'_{j\lambda}, u_{\alpha j\mu})$  in  $A$  and the matrix  $(x_{j\mu})$  defines a map  $\psi: M \rightarrow N$  such that a diagram as above commutes.  $\square$



(4.3) Proposition. Let  $(A, \underline{m})$  be an excellent local CM-ring,  $p := \text{char}(A/\underline{m})$ , and  $I_s(A)$  the ideal defining the singular locus of  $A$ . Suppose that

i) for every  $q \in \text{Reg } A$  containing  $pA$  the ring  $A_q/pA_q$  is regular,

ii) there exists a flat, reduced noetherian complete local  $A$ -algebra  $(B, \underline{n})$  such that

(ii<sub>1</sub>)  $(B, \underline{n})$  is CM and its residue field  $K$  is perfect,

(ii<sub>2</sub>) for every  $q \in \text{Reg } A$  the map  $A_q \rightarrow A_q \otimes_A B$  is regular,

(iii)  $I_s(A) \subseteq \underline{m}$ .

Then  $(A, I_s(A))$  is a CM-approximation.

Proof. Let  $q' \in \text{Spec } B$  and  $q := q' \cap A$ . If  $q \in \text{Reg } A$  then  $A_q \rightarrow B_{q'}$  is regular by ii<sub>2</sub>) and so  $q' \in \text{Reg } B$ . Thus if  $q' \not\in \text{Reg } B$  then  $q' \not\in I_s(B)$ , i.e.  $I_s(B) \supseteq I_s(A)B$ . Moreover  $I_s(B) = \sqrt{I_s(A)B}$  by Lemma (2.1).

If  $q'$  contains  $pA$  then  $A_q/pA_q$  is regular (see i)). Since  $A_q/pA_q \rightarrow B_{q'}/pB_{q'}$  is regular by base change we get  $B_{q'}/pB_{q'}$  regular too. Applying Proposition (3.7) to  $(B, \underline{n})$  we note that  $(B, I_s(B))$  is a CM-approximation. By Lemma (4.1)  $(B, I_s(A)B)$  is a CM-approximation and so  $(A, I_s(A))$  is too (see Lemma (4.2)).  $\square$

(4.4) Theorem. Let  $(A, \underline{m})$  be a reduced excellent local CM-ring,  $k := A/\underline{m}$ ,  $p := \text{char } k$  and  $I_s(A)$  the ideal defining the singular locus of  $A$ . Suppose that

i)  $[k : k^p] < \infty$  if  $p > 0$ ,

ii) for every  $q \in \text{Reg } A$  containing  $pA$  the ring  $A_q/pA_q$  is regular.

iii)  $I_s(A) \subseteq \underline{m}$ .

Then  $(A, I_s(A))$  is a CM-approximation.

Proof. If  $k$  is perfect then apply Proposition (4.3) for  $B = \hat{A}$  the completion of  $(A, \underline{m})$  (the map  $A \rightarrow \hat{A}$  is regular because  $A$  is excellent and  $\hat{A}$  is reduced because  $A$  is so).

If  $k$  is not perfect let  $K := k^{1/p^\infty}$  and  $P$  its prime subfield. Then from the following exact sequence

$$\Gamma_{K/P} = 0 \longrightarrow \Gamma_{K/k} \longrightarrow \Omega_{K/P} \otimes_K K$$

we get  $\text{rank}_K \Gamma_{K/k} \leq \text{rank}_K \Omega_{K/P} = \text{rank}_K \Omega_{K/kP} < \infty$ , where  $\Gamma_{K/k}$  denotes the imperfection module  $[M_1]$  (39.B).

Using EGA (22.2.6), or [NP] Corollary (3.6) there exists a formally smooth Noetherian complete local  $A$ -algebra  $(B, \underline{n})$  such that

1)  $B/\underline{n} \cong K$

$$2) \dim B = \dim A + \text{rank}_K \int K/k^*$$

Then the structural morphism  $A \longrightarrow B$  is regular by André-Radu Theorem (see [An], or [BR<sub>1</sub>], [BR<sub>2</sub>]) because  $A$  is excellent. Moreover  $B$  is a reduced CM-ring by [M<sub>1</sub>] (33.B). Now apply Proposition (4.3).  $\square$

(4.5) Lemma. Let  $(A, \underline{m})$  be a Noetherian henselian local ring and  $\underline{a} \subset A$  an ideal. Suppose that  $(A, \underline{a})$  is a CM-approximation. Let  $\gamma : N \longrightarrow N$  be its CM-function and  $r = \gamma(1)$ . Then a MCM  $A$ -module  $M$  is indecomposable iff  $M/\underline{a}^r M$  is indecomposable over  $A/\underline{a}^r$ .

Proof. (inspired by [Y] (2.10)). Clearly  $M/\underline{a}^r M$  is decomposable if  $M$  is so (use the Nakayama's Lemma). If  $M$  is indecomposable then  $\text{End}_A(M)$  is a local  $A$ -algebra,  $A$  being henselian. Let  $f$  be an idempotent from  $\text{End}_A(M/\underline{a}^r M)$ . Then there exists a linear  $A$ -map  $g : M \longrightarrow M$  such that  $\bar{g} := (A/\underline{a}) \otimes g = (A/\underline{a}) \otimes f$  ( $\gamma$  is a CM-function). Clearly  $\bar{g}$  is an idempotent. Since  $\text{End}_A(M)$  is local the sub- $A$ -algebra

$$\left\{ (A/\underline{a}) \otimes h \mid h \in \text{End}_A(M) \right\} \subset \text{End}_A(M/\underline{a}M)$$

is local too. Thus  $\bar{g} = 0$  or  $\bar{g} = 1$ . Then  $\underline{a} \cdot (M/\underline{a}^r M)$  contains either  $\text{Im } f$  or  $\text{Im}(1-f)$ . Since  $f$  is idempotent we get either  $\text{Im } f = \text{Im } f^r = 0$  or  $\text{Im}(1-f) = \text{Im}(1-f)^r = 0$ . Thus  $f = 0$  or  $f = 1$ .  $\square$

(4.6) Lemma. Conserving the hypothesis and the notations from (4.5), let  $M, N$  be two MCM  $A$ -modules such that  $M$  (resp.  $N$ ) is indecomposable and  $h : M \longrightarrow N$  a linear  $A$ -map. Suppose that  $(A/\underline{a}^r) \otimes_A h$  has a retraction (resp. section). Then  $h$  has a retraction (resp. section).

Proof. Since  $(A, \underline{a})$  is a CM-approximation there exists a linear  $A$ -map  $g : N \longrightarrow M$  such that  $(A/\underline{a}) \otimes g$  is a retraction (resp. section) of  $(A/\underline{a}) \otimes h$ . Then  $\text{Im}(1-gh) \subseteq \underline{a}M$  (resp.  $\text{Im}(1-hg) \subseteq \underline{a}N$ ). Since  $\text{End}_A(M)$  (resp.  $\text{End}_A(N)$ ) is a local ring we get  $gh = 1 - (1-gh)$  (resp.  $hg = 1 - (1-hg)$ ) bijective. Thus  $h$  has a retraction  $(gh)^{-1}g$  (resp. a section).  $\square$

(4.7) Let  $\underline{b}$  be an ideal in a Noetherian local ring  $(B, \underline{n})$ . Then  $\underline{b}$  is a CM-reduction ideal if the following statements hold:

- i) A MCM  $B$ -module  $M$  is indecomposable iff  $M/\underline{b}M$  is indecomposable over  $B/\underline{b}$ ,
- ii) Two indecomposable MCM  $B$ -modules  $M, N$  are isomorphic iff  $M/\underline{b}M$  and  $N/\underline{b}N$  are isomorphic over  $B/\underline{b}$ .



Note that our CM-reduction ideal is not necessarily  $\underline{n}$ -primary as in [D]. If  $\underline{b}$  is a CM-reduction ideal of  $B$  then  $\underline{b}^s$  is also one for every  $s \in \mathbb{N}$ .

(4.8) Theorem. Let  $(A, \underline{m})$  be a reduced excellent henselian local CM-ring,  $k := A/\underline{m}$ ,  $p := \text{char } k$  and  $I_s(A)$  the ideal defining the singular locus of  $A$ . Suppose that

- i)  $[k : k^p] < \infty$  if  $p > 0$ ,
- ii) for every  $q \in \text{Reg } A$  containing  $pA$  the ring  $A_q/pA_q$  is regular,
- iii)  $I_s(A) \subseteq \underline{m}$ .

Then there exists a positive integer  $r$  such that  $I_s(A)^r$  is a CM-reduction ideal of  $A$ .

The proof follows from the Lemmas (4.5), (4.6).

Let  $n_A$  be the cardinal of the isomorphism classes of indecomposable MCM  $A$ -modules.

(4.9) Corollary. Conserving the notations and hypothesis of Theorem (4.8) let  $B$  be the completion of  $A$  with respect to  $I_s(A)$ . Then

- i) A MCM  $A$ -module  $M$  is indecomposable iff  $B \otimes_A M$  is an indecomposable MCM  $B$ -module,
- ii) Two indecomposable MCM  $B$ -modules  $M, N$  are isomorphic iff  $B \otimes_A M, B \otimes_A N$  are isomorphic over  $B$ .

In particular  $n_A \leq n_B$ .

Proof. i) By Theorem (4.8) there exists  $r \in \mathbb{N}$  such that  $I_s(A)^r$  is a CM-reduction ideal of  $A$ . Let  $M$  be an indecomposable MCM  $A$ -module. Then  $B \otimes_A M$  is a MCM  $B$ -module by flatness and  $\bar{M} = M/I_s(A)^r M$  is indecomposable over  $\bar{A} := A/I_s(A)^r$ . Since  $\bar{A} \cong B/I_s(A)^r B$  it follows that  $(B \otimes_A \bar{A}) \otimes_A M$  is indecomposable over  $B \otimes_A \bar{A}$  and so  $B \otimes_A M$  is indecomposable too.

Conversely if  $B \otimes_A M$  is an indecomposable MCM  $B$ -module and  $x$  a system of parameters in  $A$  then  $x$  is a  $B \otimes_A M$ -regular sequence. Since  $A \rightarrow B$  is faithfully flat it follows that  $x$  is  $M$ -regular sequence, i.e.  $M$  is a MCM  $A$ -module. Clearly  $M$  must be indecomposable because  $B \otimes_A M$  is so.

ii) If  $B \otimes_A M \cong B \otimes_A N$  then  $B \otimes_A \bar{M} \cong B \otimes_A \bar{N}$  and so  $\bar{A} \otimes_A M \cong \bar{A} \otimes_A N$ . Thus  $M \cong N$  because  $I_s(A)^r$  is a CM-reduction ideal of  $A$ .  $\square$

(4.10) Corollary. Let  $(A, \underline{m})$  be a reduced excellent henselian local CM-ring,  $k := A/\underline{m}$ ,  $p := \text{char } k$ . Suppose that

- i)  $A$  is an isolated singularity, i.e.  $\underline{m} = I_s(A)$ ,
- ii)  $[k : k^p] < \infty$  if  $p > 0$ ,
- iii) for every  $q \in \text{Reg } A$  containing  $pA$  the ring  $A_q/pA_q$  is regular.

Then  $n_A \leq n_{\hat{A}}$ , where  $\hat{A}$  is the completion of  $(A, \underline{m})$ .

24810

Remark. When  $k$  is perfect and  $pA = 0$  then Corollary (4.10) is an easy consequence of [Y] (2.10), (2.12) and [Po] (1.3) (see our (5.6)).

(4.11) Corollary. Let  $(A, m)$  be a reduced excellent henselian local ring. Suppose that

- i)  $A$  is an isolated singularity of equal characteristic,
- ii)  $k := A/m$  is algebraically closed and  $\text{char } k \neq 2$ ,
- iii) The completion  $\hat{A}$  is a simple hypersurface singularity (resp. a singularity of type  $A_\infty, D_\infty$ ).

Then  $A$  is of finite CM-type (resp. of countable CM-type), i.e.  $A$  has a just a finite (resp. countable) set of isomorphic classes of indecomposable MCM  $A$ -modules.

The proof follows by [GK], [K], [BGS] and our (4.10).

## 5. Bounded multiplicity CM-type

(5.1). For beginning we list some definitions and facts from the Auslander-Reiten theory for the MCM modules. (see [Au<sub>3</sub>], [AR<sub>1</sub>], [Pl], [Ya], or [Y] Appendix).

Let  $(A, m)$  be a henselian local CM-ring and  $M, N$  two indecomposable MCM  $A$ -modules. A linear  $A$ -map  $f: M \rightarrow N$  is irreducible if  $f$  is not an isomorphism and given any factorization  $f = gh$  in the category  $\text{CM}(A)$ ,  $g$  has a section or  $h$  has a retraction. The AR - quiver of  $A$  is a directed graph which has as vertices the isomorphic classes of indecomposable MCM modules over  $A$  and there is an arrow from the isomorphic class of  $M$  to that of  $N$  provided there is an irreducible linear map from  $M$  to  $N$ . A chain of irreducible maps from  $M$  to  $N$  is a sequence of irreducible linear maps:  $M_0 \xrightarrow{f_1} M_1 \rightarrow \dots \xrightarrow{f_n} M_n$  with all  $M_i$  indecomposable MCM  $A$ -modules;  $n$  is called the length of the chain. If  $A$  is an isolated singularity then the AR-graph of  $A$  is locally finite, i.e. each vertex may be incident to only a finite number of other vertices (see [Au<sub>2</sub>], [AR<sub>3</sub>] and [Y] (A.18))

The following two Lemmas are just variants of [Y] Lemmas (3.1), (3.2), or [D]

§ 1.

(5.2) Lemma. (Harada-Sai lemma for MCM-modules). Let  $n$  be a positive integer,  $M_i$ ,  $0 \leq i \leq 2^n$  some indecomposable MCM  $A$ -modules and  $f_i: M_{i-1} \rightarrow M_i$ ,  $1 \leq i \leq 2^n$  some nonisomorphic linear  $A$ -maps.

Suppose that

- i)  $\underline{m}^r$  is a CM-reduction ideal of  $A$  for a certain  $r \in \mathbb{N}$ ,



ii)  $\text{length}(M_i/\underline{m}^r M_i) \leq n, \quad 0 \leq i \leq n.$

Then  $(A/\underline{m}^r) \otimes (f_{2n} \circ \dots \circ f_1) = 0.$

The proof follows easily from [HS] Lemma 12 and our Lemma (4.6).

(5.3). Lemma. Let  $n$  be a positive integer,  $M, N$  two indecomposable MCM  $A$ -modules and  $\varphi: M \rightarrow N$  a linear  $A$ -map. Suppose that

1)  $\underline{m}^r$  is a CM-reduction ideal of  $A$  for a certain  $r \in \mathbb{N}$ ,

2)  $(A/\underline{m}^r) \otimes \varphi \neq 0,$

3) there is no chain of irreducible maps from  $M$  to  $N$  of length  $< n$  which is nontrivial modulo  $\underline{m}^r$ .

Then

i) there exist a chain of irreducible maps

$$M = M_0 \xrightarrow{f_1} M_1 \rightarrow \dots \xrightarrow{f_n} M_n$$

and a linear  $A$ -map  $g: M_n \rightarrow N$  such that  $(A/\underline{m}^r) \otimes (g \circ f_n \circ \dots \circ f_1) \neq 0$

ii) there exist a chain of irreducible maps

$$N_n \xrightarrow{g_n} N_{n-1} \rightarrow \dots \xrightarrow{g_1} N_0 = N$$

and a linear  $A$ -map  $f: M \rightarrow N_n$  such that  $(A/\underline{m}^r) \otimes (g_1 \circ \dots \circ g_n \circ f) \neq 0$

... The proof follows as in [Y] (3.2).

(5.4) Theorem. Let  $(A, \underline{m})$  be a reduced excellent henselian local CM-ring,  $k := A/\underline{m}$ ,  $p = \text{char } k$ ,  $\Gamma$  the AR-quiver of  $A$  and  $\Gamma^0$  a connected component of  $\Gamma$ . Suppose that

i)  $A$  is an isolated singularity,

ii)  $\Gamma^0$  is of bounded multiplicity type, i.e. there exists  $n \in \mathbb{N}$  such that all indecomposable MCM modules  $M$  whose isomorphic classes are vertices in  $\Gamma^0$  hold  $e(M) \leq n$ .

iii)  $[k : k^p] < \infty$  if  $p > 0$ ,

iv) for every  $q \in \text{Reg } A$  containing  $pA$  the ring  $A_q/pA_q$  is regular,

Then  $\Gamma = \Gamma^0$  and  $\Gamma$  is a finite graph. In particular  $A$  is of finite CM-type.

Proof. (inspired from [Y] (3.3)). By i) we have  $I_s(A) = \underline{m}$  and it follows that  $\underline{m}^r$  is a CM-reduction ideal of  $A$  for a certain  $r \in \mathbb{N}$  (see Theorem (4.8)). Let  $x$  be a system of parameters of  $A$  and  $M$  a MCM  $A$ -module. By [M<sub>2</sub>] (14.11) we have

$$\text{length}_A(M/xM) = e(xA, M)$$

because  $x$  is a  $M$ -regular sequence. Let  $u \in \mathbb{N}$  be such that  $\underline{m}^u \subseteq xA$ . By [M<sub>2</sub>] (14.3),

(14.4) we get

$$e(xA, M) \leq e(m^u, M) = e(M)u^d$$

where  $d = \dim A$ . Choosing  $x$  in  $\underline{m}^r$  it follows

$$(1) \text{length}_A(M/\underline{m}^r M) \leq e(M)u^d$$

Let  $\mathcal{M}$  be the class of all MCM  $A$ -modules whose isomorphic classes are vertices in  $\Gamma^0$ . Using (1) we get

$$(2) \text{length}_A(M/\underline{m}^r M) \leq s := nu^d = \text{constant for every } M \in \mathcal{M}.$$

Let  $M, N$  be two indecomposable MCM  $A$ -modules and  $f: M \rightarrow N$  a linear  $A$ -map such that  $(A/\underline{m}^r) \otimes f \neq 0$ . If  $M \in \mathcal{M}$  then there is a chain of irreducible maps from  $M$  to  $N$  of length  $< t := 2^s$  which is nontrivial modulo  $\underline{m}^r$ . Otherwise there exists a chain of irreducible maps as in Lemma (5.3) i)

$$M = M_0 \xrightarrow{f_1} M_1 \rightarrow \dots \xrightarrow{f_t} M_t$$

and a linear  $A$ -map  $g: M_t \rightarrow N$  such that  $(A/\underline{m}^r) \otimes (g \circ f_t \circ \dots \circ f_1) \neq 0$ . Then  $(A/\underline{m}^r) \otimes (f_t \circ \dots \circ f_1) \neq 0$  which contradicts Lemma (5.2) ( $M_i$  are all in  $\Gamma^0$  because  $\Gamma^0$  is conex and apply (2)). In particular we get  $N \in \mathcal{M}$ . Conversely if  $N \in \mathcal{M}$  then a dual argument (using (5.3) ii) instead i)) shows that  $M \in \mathcal{M}$  and there exists a nontrivial chain of irreducible maps from  $M$  to  $N$  of length  $< t$ .

If  $M$  is a finitely generated  $A$ -module there exists a linear  $A$ -map  $f: A \rightarrow M$  such that  $(A/\underline{m}^r) \otimes f \neq 0$  (choose  $x \in M \setminus \underline{m}M$  ( $M$  is finite!) and take  $f(a) = ax$ ). If  $M \in \mathcal{M}$  then  $A \in \mathcal{M}$ . Moreover if  $M$  is an indecomposable  $A$ -module then  $M \in \mathcal{M}$  because  $A \in \mathcal{M}$ . Thus  $\Gamma^0 = \Gamma$ . Since  $\Gamma$  is locally finite and every module from  $\mathcal{M}$  can be connected with  $A$  by a chain of irreducible maps of length  $< t$  we conclude that  $\Gamma$  is finite.  $\square$

(5.5) Remark. When  $A$  is Artinian then our Theorem is a consequence of [R], [Au<sub>1</sub>]. When  $A$  is complete,  $pA = 0$  and  $k$  is perfect then our Theorem follows from [Y] (1.1).

(5.6) Remark. Another possible approach to study the CM-type is to use Artin approximation theory (see [Ar], or [Po]). Let  $(A, \underline{m})$  be a Noetherian local ring with the property of Artin approximation (shortly  $A$  is an AP-ring), i.e. for every finite system of polynomial equations  $f$  over  $A$ , every  $s \in \mathbb{N}$  and every formal solution  $\hat{y}$  of  $f$  in the completion  $\hat{A}$  of  $A$  there exists a solution  $y$  of  $f$  in  $A$  such that  $y \equiv \hat{y} \pmod{\underline{m}^s A}$ . Let  $M, N$  be two finitely generated  $A$ -modules. If  $A$  is an AP-ring then

i)  $M$  is indecomposable iff  $\hat{A} \otimes_A M$  is so,



$$\text{ii) } M \cong N \text{ iff } \hat{A} \otimes_A M \cong \hat{A} \otimes_A N.$$

For the proof note that the question can be expressed by the compatibility of some systems of polynomial equations over  $A$  (as in the proof of (4.2); but this time the equations are not linear). In particular the CM-type of  $A$  is finite if the CM-type of  $\hat{A}$  is so. Since excellent henselian local rings are AP (see [Po] Theorem (1.3)) we note that our Theorem(5.4) follows from [Y](1.1) when  $k$  is perfect and  $pA = 0$ .

## REFERENCES

- [An] M. André, Localisation de la lissité formelle, *Manuscripta Math.*, 13 (1974), 297-307.
- [Ar] M. Artin, Algebraic approximation of structures over complete local rings, *Publ. Math. IHES*, 36 (1969), 23-58.
- [Au<sub>1</sub>] M. Auslander, Representation theory of Artin algebras II, *Comm. Algebra*, 1 (1974), 269-310.
- [Au<sub>2</sub>] M. Auslander, Rational singularities and almost split sequences, *Trans. Amer. Math. Soc.*, 293 (1986), 511-531.
- [Au<sub>3</sub>] M. Auslander, Isolated singularities and existence of almost split sequences, *Proc. ICRA IV, Lect. Notes in Math.*, 1178 (1986), 194-241.
- [AR<sub>1</sub>] M. Auslander and I. Reiten, Representation theory of Artin algebras III, *Comm. Algebra*, 3 (1975), 239-294.
- [AR<sub>2</sub>] M. Auslander and I. Reiten, Representation theory of Artin algebras IV, *Comm. Algebra*, 5 (1977), 443-518.
- [BR<sub>1</sub>] A. Brezuleanu and N. Radu, Sur la localisation de la lissité formelle *C.R. Acad. Sci. Paris*, 276 (1973), 439-441.
- [BR<sub>2</sub>] A. Brezuleanu and N. Radu, Excellent rings and good separation of the module of differentials, *Rev. Roum. Math. Pures et Appl.*, 23 (1978), 1455-1470.
- [BGS] R.-O. Buchweitz, G.-M. Greuel, F. -O. Schreyer, Cohen-Macaulay modules on hypersurface singularities II, *Invent. Math.* 88 (1987), 165-183.
- [GK] G. M. Greuel and H. Knorrer, Einfache Kurvensingularitäten und torsionsfreie Moduln, *Math. Annalen* 270 (1985), 417-425.
- [D] E. Dieterich, Reduction of isolated singularities, *Comment. Math. Helvetici* 62 (1987), 654-676.
- [EGA] A. Grothendieck and J. Dieudonné, *Elements de geometrie algebrique*, IV, part 1, *Publ. Math. IHES*, 1964.
- [HS] M. Harada and Y. Sai, On categories of indecomposable modules I, *Osaka J. Math.*, 8 (1971), 309-321.
- [K] H. Knorrer, Cohen Macaulay modules on hypersurface singularities I, *Invent. Math.* 88 (1987), 153-165.
- [M<sub>1</sub>] H. Matsumura, *Commutative Algebra*, Benjamin, New York, 1970.
- [M<sub>2</sub>] H. Matsumura, *Commutative ring theory*, Cambridge University Press, 1986.
- [NP] V. Nica and D. Popescu, A structure theorem on formally smooth morphisms in positive characteristic, *J. of Algebra*, 100 (1986), 436-455.
- [P] R.S. Pierce, *Associative algebras*, Graduate Texts in Math., 88, Springer-Verlag, 1982.
- [Po] D. Popescu, General Neron desingularization and approximation, *Nagoya Math. J.*, 104 (1986), 85-115.
- [R] A. V. Roiter, Unbounded dimensionality of indecomposable representation of an algebra with an infinite number of indecomposable representations. *Izv. Akad. Nauk.*



SSSR, 32 (1968), 1275-1282 (Russian).

[S] G. Scheja and U. Storch, Lokale Verzweigungs-theorie, Schriftenreihe des Math. Inst. der Univ. Freiburg, 1974.

[Ya] K. Yamagata, On Artin rings of finite representation type, J. of Algebra, 50 (1978), 276-283.

[Y] Y. Yoshino, Brauer - Thrall type theorem for maximal Cohen-Macaulay modules, J. Math. Soc. Japan, 39 (1987), 719-739.

INCREST

Department of Mathematics  
Bd. Păcii 220, 79622 Bucharest

ROMANIA